

Dependency Pairs and Polynomial Path Orders*

Martin Avanzini and Georg Moser
 {martin.avanzini,georg.moser}@uibk.ac.at

March 2009

We show how polynomial path orders can be employed efficiently in conjunction with weak innermost dependency pairs to automatically certify polynomial runtime complexity of term rewrite systems and the polytime computability of the functions computed. The established techniques have been implemented and we provide ample experimental data to assess the new method.

Contents

1. Introduction	2
2. The Polynomial Path Order on Sequences	3
3. Complexity Analysis Based on the Dependency Pair Method	5
4. The Polynomial Path Order over Quasi-Precedences	7
5. Dependency Pairs and Polynomial Path Orders	9
6. Experimental Results	14
7. Conclusion	15
A. Appendix	15
A.1. Proof of Theorem 5.11	18
A.2. Proof of Theorem 5.14	20

* This research is partially supported by FWF (Austrian Science Fund) projects P20133.

1. Introduction

In order to measure the complexity of a (terminating) term rewrite system (TRS for short) it is natural to look at the maximal length of derivation sequences—the *derivation length*—as suggested by Hofbauer and Lautemann in [16]. More precisely, the *runtime complexity function* with respect to a (finite and terminating) TRS $\mathcal{R}$ relates the maximal derivation length to the size of the initial term, whenever the set of initial terms is restricted to constructor based terms, also called *basic* terms. The restriction to basic terms allows us to accurately express the complexity of a program through the runtime complexity of TRSs. In this paper we study and combine recent efforts for the *automatic* analysis of runtime complexities of TRSs. In [4] we introduced a restriction of the multiset path order, called *polynomial path order* (POP* for short) that induces polynomial runtime complexity if restricted to innermost rewriting. The definition of POP* employs the idea of *tiered recursion* [19]. Syntactically this amounts to a separation of arguments into *normal* and *safe* arguments, cf. [7]. Furthermore, Hirokawa and the second author introduced a variant of dependency pairs, dubbed *weak dependency pairs*, that makes the dependency pair method applicable in the context of complexity analysis, cf. [15, 14].

We show how weak innermost dependency pairs can be successfully applied in conjunction with POP*. The following example (see [11]) motivates this study. Consider the TRS $\mathcal{R}_{\text{bin}}$ encoding the function $\lambda x. \lceil \log(x+1) \rceil$ for natural numbers given as tally sequences:

$$\begin{array}{ll} 1: & \text{half}(0) \rightarrow 0 \\ 2: & \text{half}(s(0)) \rightarrow 0 \\ 3: & \text{half}(s(s(x))) \rightarrow s(\text{half}(x)) \\ 4: & \text{bits}(0) \rightarrow 0 \\ 5: & \text{bits}(s(0)) \rightarrow s(0) \\ 6: & \text{bits}(s(s(x))) \rightarrow s(\text{bits}(s(\text{half}(x)))) \end{array}$$

It is easy to see that the TRS $\mathcal{R}_{\text{bin}}$ is not compatible with POP*, even if we allow quasi-precedences, see Section 4. On the other hand, employing (weak innermost) dependency pairs, argument filtering, and the usable rules criteria in conjunction with POP*, polynomial innermost runtime complexity of $\mathcal{R}_{\text{bin}}$ can be shown fully automatically.

The combination of dependency pairs and polynomial path orders, while conceptually quite clear, turns out to be technical involved. One of the first obstacles one encounters is that the pair $(\succ_{\text{pop}*}, >_{\text{pop}*})$ cannot be used as a reduction pair in the spirit of [15], as $\succ_{\text{pop}*}$ fails to be closed under contexts. Conclusively, we start from scratch and study polynomial path orders in the context of *relative rewriting* [12]. Based on this study an incorporation of argument filterings becomes possible so that we can employ the pair $(\succ_{\text{pop}*}^{\pi}, >_{\text{pop}*}^{\pi})$ in conjunction with dependency pairs successfully. Here, $>_{\text{pop}*}^{\pi}$ refers to the order obtained by combining $>_{\text{pop}*}$ with the argument filtering π as expected, and $\succ_{\text{pop}*}^{\pi}$ denotes the extension of $>_{\text{pop}*}^{\pi}$ by term equivalence, preserving the separation of safe and normal argument positions. Note that for polynomial path orders, the integration of argument filterings is not only non-trivial, but indeed a challenging task. This is mainly due to the embodiment of tiered recursion in POP*. Thus we establish a combination of two syntactic techniques in complexity analysis. The experimental evidence given below indicates the power and in particular the efficiency of the provided results.

Our next contribution is concerned with *implicit complexity theory*, see for example [8]. A careful analysis of our main result shows that polynomial path orders in conjunction with (weak innermost) dependency pairs even induce polytime computability of the functions defined by the TRS studied. This result fits well with recent results by Marion and Péchoux on the use of restricted forms of the dependency pair method to characterise complexity classes like PTIME or PSPACE, cf. [17]. Note that both results allow to conclude, based on different restrictions, polytime computability of the functions defined by constructor TRSs, whose termination can be shown by the dependency pair method. Note that the results in [17] also capture programs admitting infeasible runtime complexities but define functions that are computable in polytime if suitable (and non-trivial) program transformations are used. Such programs are outside the scope of our results. Thus it seems that our results more directly assess the complexity of the given programs. Note that our tool provides (for the first time) a fully automatic application of the dependency pair method in the context of implicit complexity theory.¹

The rest of the paper is organised as follows. In Section 2 we present basic notions and recall (briefly) the *path order for FP* from [1]. We then briefly recall dependency pairs in the context of complexity analysis from [15, 14], cf. Section 3. In Section 4 we present polynomial path orders over quasi-precedences. Our main results are presented in Section 5. We continue with experimental results in Section 6, and conclude in Section 7.

2. The Polynomial Path Order on Sequences

We assume familiarity with the basics of term rewriting, see [6, 20]. Let $\mathcal{V}$ denote a countably infinite set of variables and $\mathcal{F}$ a signature, containing at least one constant. The set of terms over $\mathcal{F}$ and $\mathcal{V}$ is denoted as $\mathcal{T}(\mathcal{F}, \mathcal{V})$ and the set of ground terms as $\mathcal{T}(\mathcal{F})$. We write $\mathbf{Fun}(t)$ and $\mathbf{Var}(t)$ for the set of function symbols and variables appearing in t , respectively. The root symbol $\mathbf{rt}(t)$ of a term t is defined as usual and the (proper) subterm relation is denoted as $\sqsubseteq$ ($\triangleleft$). We write $s|_p$ for the *subterm* of s at position p . The *size* $|t|$ of a term t is defined as usual and the *width* of t is defined as $\mathbf{width}(t) := \max\{n, \mathbf{width}(t_1), \dots, \mathbf{width}(t_n)\}$ if $t = f(t_1, \dots, t_n)$ and $n > 0$ or $\mathbf{width}(t) = 1$ else. Let $\succsim$ be a preorder on the signature $\mathcal{F}$, called *quasi-precedence* or simply *precedence*. Based on $\succsim$ we define an equivalence $\approx$ on terms: $s \approx t$ if either (i) $s = t$ or (ii) $s = f(s_1, \dots, s_n)$, $t = g(t_1, \dots, t_n)$, $f \approx g$ and there exists a permutation π such that $s_i \approx t_{\pi(i)}$. For a preorder $\succsim$, we use $\succsim^{\text{mul}}$ for the multiset extension of $\succsim$, which is again a preorder. The proper order (equivalence) induced by $\succsim^{\text{mul}}$ is written as $\succ^{\text{mul}}$ ($\approx^{\text{mul}}$).

A *term rewrite system* (TRS for short) $\mathcal{R}$ over $\mathcal{T}(\mathcal{F}, \mathcal{V})$ is a *finite* set of rewrite rules $l \rightarrow r$, such that $l \notin \mathcal{V}$ and $\mathbf{Var}(l) \supseteq \mathbf{Var}(r)$. We write $\rightarrow_{\mathcal{R}}$ ($\stackrel{i}{\rightarrow}_{\mathcal{R}}$) for the induced (innermost) rewrite relation. The set of defined function symbols is denoted as $\mathcal{D}$, while the constructor symbols are collected in $\mathcal{C}$, clearly $\mathcal{F} = \mathcal{D} \cup \mathcal{C}$. We use $\mathbf{NF}(\mathcal{R})$ to denote the set of normal forms of $\mathcal{R}$ and set $\mathbf{Val} := \mathcal{T}(\mathcal{C}, \mathcal{V})$, the elements of $\mathbf{Val}$ are called

¹ In this context it is perhaps interesting to note that for a variant of the TRS $\mathcal{R}_{\text{bin}}$, studied in [17], our tool verifies polytime computability fully automatically. See also [5] for the description of a small tool that implements related characterisations of the class of polynomial time computable functions.

values. A TRS is called *completely defined* if normal forms coincide with values. We define $\mathcal{T}_b := \{f(v_1, \dots, v_n) \mid f \in \mathcal{D} \text{ and } v_i \in \text{Val}\}$ as the set of *basic terms*. A TRS $\mathcal{R}$ is a *constructor TRS* if $l \in \mathcal{T}_b$ for all $l \rightarrow r \in \mathcal{R}$. Let $\mathcal{Q}$ denote a TRS. The *generalised restricted rewrite relation* $\xrightarrow{\mathcal{Q}}_{\mathcal{R}}$ is the restriction of $\rightarrow_{\mathcal{R}}$ where all arguments of the redex are in normal form with respect to the TRS $\mathcal{Q}$ (see [21]). We define the (innermost) relative rewriting relation (denoted as $\xrightarrow{\cdot}_{\mathcal{R}/\mathcal{S}}$) as follows:

$$\xrightarrow{\cdot}_{\mathcal{R}/\mathcal{S}} := \xrightarrow{\mathcal{R} \cup \mathcal{S}}_{\mathcal{S}}^* \cdot \xrightarrow{\mathcal{R} \cup \mathcal{S}}_{\mathcal{R}} \cdot \xrightarrow{\mathcal{R} \cup \mathcal{S}}_{\mathcal{S}}^*.$$

Similarly, we set $\xrightarrow{\cdot}_{\mathcal{R}/\mathcal{S}}^{\varepsilon} := \xrightarrow{\mathcal{R} \cup \mathcal{S}}_{\mathcal{S}}^* \cdot \xrightarrow{\mathcal{R} \cup \mathcal{S}}_{\mathcal{R}}^{\varepsilon} \cdot \xrightarrow{\mathcal{R} \cup \mathcal{S}}_{\mathcal{S}}^*$, to define an (*innermost*) *relative root-step*.

A *polynomial interpretation* is a well-founded and monotone algebra $(\mathcal{A}, >)$ with carrier $\mathbb{N}$ such that $>$ is the usual order on natural numbers and all interpretation functions $f_{\mathcal{A}}$ are polynomials. Let $\alpha: \mathcal{V} \rightarrow \mathcal{A}$ denote an *assignment*, then we write $[\alpha]_{\mathcal{A}}(t)$ for the evaluation of term t with respect to $\mathcal{A}$ and α . A polynomial interpretation is called a *strongly linear interpretation* (SLI for short) if all function symbols are interpreted by *weight functions* $f_{\mathcal{A}}(x_1, \dots, x_n) = \sum_{i=1}^n x_i + c$ with $c \in \mathbb{N}$. The *derivation length* of a terminating term s with respect to $\rightarrow$ is defined as $\text{dl}(s, \rightarrow) := \max\{n \mid \exists t. s \rightarrow^n t\}$, where $\rightarrow^n$ denotes the n -fold application of $\rightarrow$. The *innermost runtime complexity function* $\text{rc}_{\mathcal{R}}^i$ with respect to a TRS $\mathcal{R}$ is defined as $\text{rc}_{\mathcal{R}}^i(n) := \max\{\text{dl}(t, \xrightarrow{\cdot}_{\mathcal{R}}) \mid t \in \mathcal{T}_b \text{ and } |t| \leq n\}$. If no confusion can arise $\text{rc}_{\mathcal{R}}^i$ is simply called *runtime complexity function*.

Below we recall the bare essentials of the polynomial path order $\blacktriangleright$ on sequences (POP for short) as put forward in [1]. We kindly refer the reader to [1, 4] for motivation and examples. We recall the definition of *finite approximations* $\blacktriangleright_k^l$ of $\blacktriangleright$. The latter is conceived as the *limit* of these approximations. The domain of this order are so called *sequences* $\text{Seq}(\mathcal{F}, \mathcal{V}) := \mathcal{T}(\mathcal{F} \cup \{\circ\}, \mathcal{V})$. Here $\mathcal{F}$ is a finite signature and $\circ \notin \mathcal{F}$ a fresh variadic function symbol, used to form sequences. We denote sequences $\circ(s_1, \dots, s_n)$ by $[s_1 \cdots s_n]$ and write $a :: [b_1 \cdots b_n]$ for the sequence $[a \ b_1 \cdots b_n]$.

Let $\succsim$ denote a precedence. The order $\blacktriangleright_k^l$ is based on an auxiliary order $\succ_k^l$ (and the equivalence $\approx$ on terms defined above). Below we set $\gtrsim_k^l := \succ_k^l \cup \approx$. We write $\{\{t_1, \dots, t_n\}\}$ to denote multisets and $\uplus$ for the multiset sum.

Definition 2.1. Let $k, l \geq 1$. The order $\succ_k^l$ induced by $\succsim$ is inductively defined as follows: $s \succ_k^l t$ for $s = f(s_1, \dots, s_n)$ or $s = [s_1 \cdots s_n]$ if either

- (i) $s_i \gtrsim_k^l t$ for some $i \in \{1, \dots, n\}$, or
- (ii) $s = f(s_1, \dots, s_n)$, $t = g(t_1, \dots, t_m)$ with $f \succ g$ or $t = [t_1 \cdots t_m]$, $s \succ_k^{l-1} t_j$ for all $j \in \{1, \dots, m\}$, and $m < k + \text{width}(s)$,
- (iii) $s = [s_1 \cdots s_n]$, $t = [t_1 \cdots t_m]$ and the following properties hold:
 - $\{\{t_1, \dots, t_m\}\} = N_1 \uplus \cdots \uplus N_n$ for some multisets $N_1, \dots, N_n$, and
 - there exists $i \in \{1, \dots, n\}$ such that $\{\{s_i\}\} \not\approx^{\text{mul}} N_i$, and
 - for all $1 \leq i \leq n$ such that $\{\{s_i\}\} \approx^{\text{mul}} N_i$ we have $s_i \succ_k^l r$ for all $r \in N_i$, and $m < k + \text{width}(s)$.

Definition 2.2. Let $k, l \geq 1$. The *approximation* $\blacktriangleright_k^l$ of the polynomial path order on sequences induced by $\succsim$ is inductively defined as follows: $s \blacktriangleright_k^l t$ for $s = f(s_1, \dots, s_n)$ or $s = [s_1 \cdots s_n]$ if either $s \succ_k^l t$ or

- (i) $s_i \blacktriangleright_k^l t$ for some $i \in \{1, \dots, n\}$,
- (ii) $s = f(s_1, \dots, s_n)$, $t = [t_1 \cdots t_m]$, and the following properties hold:
 - $s \blacktriangleright_k^{l-1} t_{j_0}$ for some $j_0 \in \{1, \dots, m\}$,
 - $s \succ_k^{l-1} t_j$ for all $j \neq j_0$, and $m < k + \text{width}(s)$,
- (iii) $s = f(s_1, \dots, s_n)$, $t = g(t_1, \dots, t_m)$, $f \sim g$ and $[s_1 \cdots s_n] \blacktriangleright_k^l [t_1 \cdots t_m]$, or
- (iv) $s = [s_1 \cdots s_n]$, $t = [t_1 \cdots t_m]$ and the following properties hold:
 - $\{\{t_1, \dots, t_m\}\} = N_1 \uplus \cdots \uplus N_n$ for some multisets $N_1, \dots, N_n$, and
 - there exists $i \in \{1, \dots, n\}$ such that $\{\{s_i\}\} \not\approx^{\text{mul}} N_i$, and
 - for all $1 \leq i \leq n$ such that $\{\{s_i\}\} \approx^{\text{mul}} N_i$ we have $s_i \blacktriangleright_k^l r$ for all $r \in N_i$, and $m < k + \text{width}(s)$.

Above we set $\blacktriangleright_k^l := \blacktriangleright_k^l \cup \approx$ and abbreviate $\blacktriangleright_k^k$ as $\blacktriangleright_k$ in the following. Note that the empty sequence is minimal with respect to both orders. It is easy to see that for $k \leq l$, we have $\succ_k \subseteq \succ_l$ and $\blacktriangleright_k \subseteq \blacktriangleright_l$. Note that $s \blacktriangleright_k t$ implies that $\text{width}(t) < \text{width}(s) + k$. For a fixed approximation $\blacktriangleright_k$, we define the length of its longest decent as follows: $G_k(t) := \max\{n \mid t = t_0 \blacktriangleright_k t_1 \blacktriangleright_k \cdots \blacktriangleright_k t_n\}$. The following proposition is a reformulation of [1, Lemma 6].

Proposition 2.3. *Let $k \in \mathbb{N}$. There exists a polynomial interpretation $\mathcal{A}$ such that $G_k(t) \leq [\alpha]_{\mathcal{A}}(t)$ for all assignments $\alpha: \mathcal{V} \rightarrow \mathbb{N}$. As a consequence, for all terms $f(t_1, \dots, t_n)$ with $[\alpha]_{\mathcal{A}}(t_i) = O(|t_i|)$, $G_k(f(t_1, \dots, t_n))$ is bounded by a polynomial p in the size of t , where p depends on k only.*

Observe that the polynomial interpretation $\mathcal{A}$ as employed in the proposition fulfils: $\circ_{\mathcal{A}}(m_1, \dots, m_n) = \sum_{i=1}^n m_i + n$. In particular, we have $[\alpha]_{\mathcal{A}}(\square) = 0$.

3. Complexity Analysis Based on the Dependency Pair Method

In this section, we briefly recall the central definitions and results established in [15, 14]. We kindly refer the reader to [15, 14] for further examples and underlying intuitions. Let $\mathcal{X}$ be a set of symbols. We write $C\langle t_1, \dots, t_n \rangle_{\mathcal{X}}$ to denote $C[t_1, \dots, t_n]$, whenever $\text{rt}(t_i) \in \mathcal{X}$ for all $i \in \{1, \dots, n\}$ and C is a n -hole context containing no symbols from $\mathcal{X}$. We set $\mathcal{D}^{\sharp} := \mathcal{D} \cup \{f^{\sharp} \mid f \in \mathcal{D}\}$ with each $f^{\sharp}$ a fresh function symbol. Further, for $t = f(t_1, \dots, t_n)$ with $f \in \mathcal{D}$, we set $t^{\sharp} := f^{\sharp}(t_1, \dots, t_n)$.

Definition 3.1. Let $\mathcal{R}$ be a TRS. If $l \rightarrow r \in \mathcal{R}$ and $r = C\langle u_1, \dots, u_n \rangle_{\mathcal{D}}$ then $l^\# \rightarrow \text{COM}(u_1^\#, \dots, u_n^\#)$ is called a *weak innermost dependency pair* of $\mathcal{R}$. Here $\text{COM}(t) = t$ and $\text{COM}(t_1, \dots, t_n) = \mathbf{c}(t_1, \dots, t_n)$, $n \neq 1$, for a fresh constructor symbol $\mathbf{c}$, the *compound symbol*. The set of all weak innermost dependency pairs is denoted by $\text{WIDP}(\mathcal{R})$.

Example 3.2. Reconsider the example $\mathcal{R}_{\text{bits}}$ from the introduction. The set of weak innermost dependency pairs $\text{WIDP}(\mathcal{R}_{\text{bits}})$ is given by

$$\begin{array}{ll} 7: & \text{half}^\#(0) \rightarrow \mathbf{c}_1 & 10: & \text{bits}^\#(0) \rightarrow \mathbf{c}_3 \\ 8: & \text{half}^\#(\mathbf{s}(0)) \rightarrow \mathbf{c}_2 & 11: & \text{bits}^\#(\mathbf{s}(0)) \rightarrow \mathbf{c}_4 \\ 9: & \text{half}^\#(\mathbf{s}(\mathbf{s}(x))) \rightarrow \text{half}^\#(x) & 12: & \text{bits}^\#(\mathbf{s}(\mathbf{s}(x))) \rightarrow \text{bits}^\#(\mathbf{s}(\text{half}(x))) \end{array}$$

We write $f \triangleright_{\mathbf{d}} g$ if there exists a rewrite rule $l \rightarrow r \in \mathcal{R}$ such that $f = \text{rt}(l)$ and g is a defined symbol in $\text{Fun}(r)$. For a set $\mathcal{G}$ of defined symbols we denote by $\mathcal{R}|_{\mathcal{G}}$ the set of rewrite rules $l \rightarrow r \in \mathcal{R}$ with $\text{rt}(l) \in \mathcal{G}$. The set $\mathcal{U}(t)$ of usable rules of a term t is defined as $\mathcal{R}|_{\{g \mid f \triangleright_{\mathbf{d}}^* g \text{ for some } f \in \text{Fun}(t)\}}$. Finally, we define $\mathcal{U}(\mathcal{P}) = \bigcup_{l \rightarrow r \in \mathcal{P}} \mathcal{U}(r)$.

Example 3.3 (Example 3.2 continued). The usable rules of $\text{WIDP}(\mathcal{R}_{\text{bits}})$ consist of the following rules: 1: $\text{half}(0) \rightarrow 0$, 2: $\text{half}(\mathbf{s}(0)) \rightarrow 0$, and 3: $\text{half}(\mathbf{s}(\mathbf{s}(x))) \rightarrow \text{half}(x)$.

The following proposition allows the analysis of the (innermost) runtime complexity through the study of (innermost) relative rewriting, see [15] for the proof.

Proposition 3.4. Let $\mathcal{R}$ be a TRS, let t be a basic terminating term, and let $\mathcal{P} = \text{WIDP}(\mathcal{R})$. Then $\text{dl}(t, \xrightarrow{\cdot}_{\mathcal{R}}) \leq \text{dl}(t^\#, \xrightarrow{\cdot}_{\mathcal{U}(\mathcal{P}) \cup \mathcal{P}})$. Moreover, if $\mathcal{P}$ is non-duplicating and $\mathcal{U}(\mathcal{P}) \subseteq >_{\mathcal{A}}$ for some SLI $\mathcal{A}$. Then there exist constants $K, L \geq 0$ (depending on $\mathcal{P}$ and $\mathcal{A}$ only) such that $\text{dl}(t, \xrightarrow{\cdot}_{\mathcal{R}}) \leq K \cdot \text{dl}(t^\#, \xrightarrow{\cdot}_{\mathcal{P}/\mathcal{U}(\mathcal{P})}) + L \cdot |t^\#|$.

This approach admits also an integration of *dependency graphs* [2] in the context of complexity analysis. The nodes of the *weak innermost dependency graph* $\text{WIDG}(\mathcal{R})$ are the elements of $\mathcal{P}$ and there is an arrow from $s \rightarrow t$ to $u \rightarrow v$ if there exist a context C and substitutions σ, τ such that $t\sigma \xrightarrow{\cdot}_{\mathcal{R}}^* C[u\tau]$. Let $\mathcal{G} = \text{WIDG}(\mathcal{R})$; a *strongly connected component* (SCC for short) in $\mathcal{G}$ is a maximal *strongly connected subgraph*. We write $\mathcal{G}/\equiv$ for the *congruence graph*, where $\equiv$ is the equivalence relation induced by SCCs.

Example 3.5 (Example 3.2 continued). $\mathcal{G} = \text{WIDG}(\mathcal{R}_{\text{bits}})$ consists of the nodes (7)–(12) as mentioned in Example 3.2 and has the following shape:

$$\begin{array}{ccc} & \cap & \\ 7 & \longleftarrow 9 & \longrightarrow 8 \\ & \cap & \end{array} \quad \begin{array}{ccc} & \cap & \\ 10 & & 12 & \longrightarrow 11 \\ & \cap & \end{array}$$

The only non-trivial SCCs in $\mathcal{G}$ are $\{9\}$ and $\{12\}$. Hence $\mathcal{G}/\equiv$ consists of the nodes $[7]_{\equiv}$ – $[12]_{\equiv}$, and edges $([a]_{\equiv}, [b]_{\equiv})$ for edges (a, b) in $\mathcal{G}$. Here $[a]_{\equiv}$ denotes the equivalence class of a .

We set $L(t) := \max\{\text{dl}(t, \xrightarrow{\mathcal{P}_m/\mathcal{S}}) \mid (\mathcal{P}_1, \dots, \mathcal{P}_m) \text{ a path in } \mathcal{G}/\equiv, \mathcal{P}_1 \in \text{Src}\}$, where Src denote the set of source nodes from $\mathcal{G}/\equiv$ and $\mathcal{S} = \mathcal{P}_1 \cup \dots \cup \mathcal{P}_{m-1} \cup \mathcal{U}(\mathcal{P}_1 \cup \dots \cup \mathcal{P}_m)$. The proposition allows the use of different techniques to analyse polynomial runtime complexity on separate paths, cf. [14].

Proposition 3.6. *Let $\mathcal{R}$, $\mathcal{P}$, and t be as above. Then there exists a polynomial p (depending only on $\mathcal{R}$) such that $\text{dl}(t^\sharp, \xrightarrow{\mathcal{P}/\mathcal{U}(\mathcal{P})}) \leq p(L(t^\sharp))$.*

4. The Polynomial Path Order over Quasi-Precedences

In this section, we briefly recall the central definitions and results established in [4, 5] on the *polynomial path order*. We employ the variant of POP* based on quasi-precedences, cf. [5].

As mentioned in the introduction, POP* relies on tiered recursion, which is captured by the notion of *safe mapping*. A *safe mapping* **safe** is a function that associates with every n -ary function symbol f the set of *safe argument positions*. If $f \in \mathcal{D}$ then $\text{safe}(f) \subseteq \{1, \dots, n\}$, for $f \in \mathcal{C}$ we fix $\text{safe}(f) = \{1, \dots, n\}$. The argument positions not included in $\text{safe}(f)$ are called *normal* and denoted by $\text{nrm}(f)$. We extend **safe** to terms $t \notin \mathcal{V}$ as follows: we define $\text{safe}(f(t_1, \dots, t_n)) := \{t_{i_1}, \dots, t_{i_p}\}$ where $\text{safe}(f) = \{i_1, \dots, i_p\}$, likewise we define $\text{nrm}(f(t_1, \dots, t_n)) := \{t_{j_1}, \dots, t_{j_q}\}$ where $\text{nrm}(f) = \{j_1, \dots, j_q\}$. Not every precedence is suitable for $\succ_{\text{pop}^*}$, in particular we need to assert that constructors are minimal.

We say that a precedence $\succsim$ is *admissible* for POP* if the following is satisfied: (i) $f \succ g$ with $g \in \mathcal{D}$ implies $f \in \mathcal{D}$, and (ii) if $f \approx g$ then $f \in \mathcal{D}$ if and only if $g \in \mathcal{D}$. In the sequel we assume any precedence is admissible. We extend the equivalence $\approx$ to the context of safe mapping: $s \approx^{\text{safe}} t$, if (i) $s = t$, or (ii) $s = f(s_1, \dots, s_n)$, $t = g(t_1, \dots, t_n)$, $f \approx g$ and there exists a permutation π so that $s_i \approx^{\text{safe}} t_{\pi(i)}$, where $i \in \text{safe}(f)$ if and only if $\pi(i) \in \text{safe}(g)$ for all $i \in \{1, \dots, n\}$. Similar to POP, the definition of the polynomial path order $\succ_{\text{pop}^*}$ makes use of an auxiliary order $\succ_{\text{pop}}$.

Definition 4.1. The auxiliary order $\succ_{\text{pop}}$ induced by $\succsim$ and **safe** is inductively defined as follows: $s = f(s_1, \dots, s_n) \succ_{\text{pop}} t$ if either

- (i) $s_i \succ_{\text{pop}} t$ for some $i \in \{1, \dots, n\}$, and if $f \in \mathcal{D}$ then $i \in \text{nrm}(f)$, or
- (ii) $t = g(t_1, \dots, t_m)$, $f \succ g$, $f \in \mathcal{D}$ and $s \succ_{\text{pop}} t_j$ for all $j \in \{1, \dots, m\}$.

Definition 4.2. The *polynomial path order* $\succ_{\text{pop}^*}$ induced by $\succsim$ and **safe** is inductively defined as follows: $s = f(s_1, \dots, s_n) \succ_{\text{pop}^*} t$ if either $s \succ_{\text{pop}} t$ or

- (i) $s_i \succ_{\text{pop}^*} t$ for some $i \in \{1, \dots, n\}$, or
- (ii) $t = g(t_1, \dots, t_m)$, $f \succ g$, $f \in \mathcal{D}$, and
 - $s \succ_{\text{pop}^*} t_{j_0}$ for some $j_0 \in \text{safe}(g)$, and
 - for all $j \neq j_0$ either $s \succ_{\text{pop}} t_j$, or $s \triangleright t_j$ and $j \in \text{safe}(g)$, or

(iii) $t = g(t_1, \dots, t_m)$, $f \approx g$, $\text{nrm}(s) >_{\text{pop}^*}^{\text{mul}} \text{nrm}(t)$ and $\text{safe}(s) \succcurlyeq_{\text{pop}^*}^{\text{mul}} \text{safe}(t)$.

Above we set $\succcurlyeq_{\text{pop}} := >_{\text{pop}} \cup \approx^{\text{safe}}$ and $\succcurlyeq_{\text{pop}^*} := >_{\text{pop}^*} \cup \approx^{\text{safe}}$ below. Here $>_{\text{pop}^*}^{\text{mul}}$ and $\succcurlyeq_{\text{pop}^*}^{\text{mul}}$ refer to the strict and weak multiset extension of $\succcurlyeq_{\text{pop}^*}$ respectively.

The intuition of $>_{\text{pop}}$ is to deny any recursive call, whereas $>_{\text{pop}^*}$ allows predicative recursion: by the restrictions imposed by **safe**, recursion needs to be performed on normal arguments, while a recursively computed result must only be used in a safe argument position, compare [7]. Note that the alternative $s \triangleright t_j$ for $j \in \text{safe}(g)$ in Definition 4.2(ii) guarantees that POP^* characterises the class of polytime computable functions, cf. [4]. The proof of the next theorem follows the pattern of the proof of main theorem in [4], but the result is stronger due to the extension to quasi-precedences.

Theorem 4.3. *Let $\mathcal{R}$ be a constructor TRS. If $\mathcal{R}$ is compatible with $>_{\text{pop}^*}$, i.e., $\mathcal{R} \subseteq >_{\text{pop}^*}$, then the innermost runtime complexity $\text{rc}_{\mathcal{R}}^i$ induced is polynomially bounded.*

Note that Theorem 4.3 is too weak to handle the TRS $\mathcal{R}_{\text{bits}}$ as the (necessary) restriction to an admissible precedence is too strong. To rectify this, we suit POP^* so that it can be used in conjunction with weak (innermost) dependency pairs.

An argument filtering (for a signature $\mathcal{F}$) is a mapping π that assigns to every n -ary function symbol $f \in \mathcal{F}$ an argument position $i \in \{1, \dots, n\}$ or a (possibly empty) list $\{i_1, \dots, i_m\}$ of argument positions with $1 \leq i_1 < \dots < i_m \leq n$. The signature $\mathcal{F}_\pi$ consists of all function symbols f such that $\pi(f)$ is some list $\{i_1, \dots, i_m\}$, where in $\mathcal{F}_\pi$ the arity of f is m . Every argument filtering π induces a mapping from $\mathcal{T}(\mathcal{F}, \mathcal{V})$ to $\mathcal{T}(\mathcal{F}_\pi, \mathcal{V})$, also denoted by π :

$$\pi(t) = \begin{cases} t & \text{if } t \text{ is a variable} \\ \pi(t_i) & \text{if } t = f(t_1, \dots, t_n) \text{ and } \pi(f) = i \\ f(\pi(t_{k_1}), \dots, \pi(t_{k_m})) & \text{if } t = f(t_1, \dots, t_n) \text{ and } \pi(f) = \{i_1, \dots, i_m\} \end{cases}.$$

Definition 4.4. Let π denote an argument filtering, and $>_{\text{pop}^*}$ a polynomial path order. We define $s >_{\text{pop}^*}^\pi t$ if and only if $\pi(s) >_{\text{pop}^*} \pi(t)$, and likewise $s \succcurlyeq_{\text{pop}^*}^\pi t$ if and only if $\pi(s) \succcurlyeq_{\text{pop}^*} \pi(t)$.

Example 4.5 (Example 3.2 continued). Let π be defined as follows: $\pi(\text{half}) = 1$ and $\pi(f) = \{1, \dots, n\}$ for each n -ary function symbol other than **half**. Compatibility of $\text{WIDP}(\mathcal{R}_{\text{bits}})$ with $>_{\text{pop}^*}^\pi$ amounts to the following set of order constraints:

$$\begin{array}{lll} \text{half}^\sharp(0) >_{\text{pop}^*} c_1 & \text{bits}^\sharp(0) >_{\text{pop}^*} c_3 & \text{half}^\sharp(s(s(x))) >_{\text{pop}^*} \text{half}^\sharp(x) \\ \text{half}^\sharp(s(0)) >_{\text{pop}^*} c_2 & \text{bits}^\sharp(s(0)) >_{\text{pop}^*} c_4 & \text{bits}^\sharp(s(s(x))) >_{\text{pop}^*} \text{bits}^\sharp(s(x)) \end{array}$$

In order to define a POP^* instance $>_{\text{pop}^*}$, we set $\text{safe}(\text{bits}^\sharp) = \text{safe}(\text{half}) = \text{safe}(\text{half}^\sharp) = \emptyset$ and $\text{safe}(s) = \{1\}$. Furthermore, we define an (admissible) precedence: $0 \approx c_1 \approx c_2 \approx c_3 \approx c_4$. The easy verification of $\text{WIDP}(\mathcal{R}_{\text{bits}}) \subseteq >_{\text{pop}^*}^\pi$ is left to the reader.

5. Dependency Pairs and Polynomial Path Orders

Motivated by Example 4.5, we show in this section that the pair $(\succsim_{\text{pop}^*}^\pi, >_{\text{pop}^*}^\pi)$ can play the role of a *safe* reduction pair, cf. [15, 14]. Let $\mathcal{R}$ be a TRS over a signature $\mathcal{F}$ that is innermost terminating. In the sequel $\mathcal{R}$ is kept fixed. Moreover, we fix some safe mapping **safe**, an admissible precedence $\succsim$, and an argument filtering π . We refer to the induced POP* instance by $>_{\text{pop}^*}^\pi$.

We adapt **safe** to $\mathcal{F}_\pi$ in the obvious way: for each $f_\pi \in \mathcal{F}_\pi$ with corresponding $f \in \mathcal{F}$, we define $\text{safe}(f_\pi) := \text{safe}(f) \cap \pi(f)$, and likewise $\text{nrm}(f_\pi) := \text{nrm}(f) \cap \pi(f)$. Set $\text{Val}_\pi := \mathcal{T}(\mathcal{C}_\pi, \mathcal{V})$. Based on $\mathcal{F}_\pi$ we define the *normalised signature* $\mathcal{F}_\pi^n := \{f^n \mid f \in \mathcal{F}_\pi\}$ where the arity of f^n is $|\text{nrm}(f)|$. We extend $\succsim$ to $\mathcal{F}_\pi^n$ by $f^n \succsim g^n$ if and only if $f \succsim g$. Let $\mathbf{s}$ be a fresh constant that is minimal with respect to $\succsim$. We introduce the *Buchholz norm* of t (denoted as $\|t\|$) a term complexity measure that fits well with the definition of POP*. Set $\|t\| := 1 + \max\{n, \|t_1\|, \dots, \|t_n\|\}$ for $t = f(t_1, \dots, t_n)$ and $\|t\| := 1$, otherwise. In the following we define an embedding from the relative rewriting relation $\xrightarrow[\mathcal{R}/\mathcal{S}]{\varepsilon}$ into $\blacktriangleright_k$, such that k depends only on TRSs $\mathcal{R}$ and $\mathcal{S}$. This embedding provides the technical tool to measure the number of root steps in a given derivation through the number of descent in $\blacktriangleright_k$. Hence Proposition 2.3 becomes applicable to establishing our main result. This intuition is cast into the next definition.

Definition 5.1. A *predicative interpretation* is a pair of mappings (S_π, N_π) from terms to sequences $\text{Seq}(\mathcal{F}_\pi^n \cup \{\mathbf{s}\}, \mathcal{V})$ defined as follows. We assume for $\pi(t) = f(\pi(t_1), \dots, \pi(t_n))$ that $\text{safe}(f) = \{i_1, \dots, i_p\}$ and $\text{nrm}(f) = \{j_1, \dots, j_q\}$.

$$S_\pi(t) := \begin{cases} [] & \text{if } \pi(t) \in \text{Val}_\pi, \\ [f^n(N_\pi(t_{j_1}), \dots, N_\pi(t_{j_q})) S_\pi(t_{i_1}) \cdots S_\pi(t_{i_p})] & \text{if } \pi(t) \notin \text{Val}_\pi. \end{cases}$$

$$N_\pi(t) := S_\pi(t) :: \text{BN}_\pi(t)$$

Here the function BN_π maps a term t to the sequence $[\mathbf{s} \cdots \mathbf{s}]$ with $\|\pi(t)\|$ occurrences of the constant $\mathbf{s}$.

Note that as a direct consequence of the definitions we obtain $\text{width}(N_\pi(t)) = \|\pi(t)\| + 1$ for all terms t .

Lemma 5.2. *There exists a polynomial p such that $G_k(N_\pi(t)) \leq p(|t|)$ for every basic term t . The polynomial p depends only on k .*

Proof. Suppose $t = f(v_1, \dots, v_n)$ is a basic term with $\text{safe}(f) = \{i_1, \dots, i_p\}$ and $\text{nrm}(f) = \{j_1, \dots, j_q\}$. The only non-trivial case is when $\pi(t) \notin \text{Val}_\pi$. Then

$$N_\pi(t) = [u S_\pi(v_{i_1}) \cdots S_\pi(v_{i_p})] :: \text{BN}_\pi(t)$$

where $u = f^n(N_\pi(v_{j_1}), \dots, N_\pi(v_{j_q}))$. Note that $S_\pi(v_i) = []$ for $i \in \{i_1, \dots, i_p\}$. Let $\mathcal{A}$ denote a polynomial interpretation fulfilling Proposition 2.3. Using the assumption $\circ_{\mathcal{A}}(m_1, \dots, m_n) = \sum_{i=1}^n m_i + n$, it is easy to see that $G_k(N_\pi(t))$ is bounded linear in $\|\pi(t)\| \leq |t|$ and $[\alpha]_{\mathcal{A}}(u)$. As $N_\pi(v_j) = [] \mathbf{s} \cdots \mathbf{s}$ with $\|\pi(v_j)\| \leq |t|$ occurrences of $\mathbf{s}$, $G_k(N_\pi(v_j))$ is linear in $|t|$. Hence from Proposition 2.3 we conclude that $G_k(N_\pi(t))$ is polynomially bounded in $|t|$. $\square$

The next sequence of lemmas shows that the relative rewriting relation $\xrightarrow{\varepsilon}_{\mathcal{R}/S}$ is embeddable into $\blacktriangleright_k$.

Lemma 5.3. *Suppose $s >_{\text{pop}^*}^\pi t$ such that $\pi(s\sigma) \in \text{Val}_\pi$. Then $S_\pi(s\sigma) = [] = S_\pi(t\sigma)$ and $N_\pi(s\sigma) \blacktriangleright_1 N_\pi(t\sigma)$.*

Proof. Let $\pi(s\sigma) \in \text{Val}_\pi$, and suppose $s >_{\text{pop}^*}^\pi t$, i.e., $\pi(s) >_{\text{pop}^*} \pi(t)$ holds. Observe that since $\pi(s) \in \text{Val}_\pi$ and due to our assumptions on safe mappings, only clause (1) from the definition of $>_{\text{pop}^*}$ (or respectively $>_{\text{pop}}$) is applicable. And thus $\pi(t)$ is a subterm of $\pi(s)$ modulo the equivalence $\approx$. We conclude $\pi(t\sigma) \in \text{Val}_\pi$, and hence $S_\pi(s\sigma) = [] = S_\pi(t\sigma)$. Finally, notice that $\|\pi(s\sigma)\| > \|\pi(t\sigma)\|$ as $\pi(t\sigma)$ is a subterm of $\pi(s\sigma)$. Thus $N_\pi(s\sigma) \blacktriangleright_1 N_\pi(t\sigma)$ follows as well. $\square$

To improve the clarity of the exposition, we concentrate on the curcial cases in the proofs of the following lemma. The interested reader is kindly referred to [3] for the full proof.

Lemma 5.4. *Suppose $s >_{\text{pop}}^\pi t$ such that $\pi(s\sigma) = f(\pi(s_1\sigma), \dots, \pi(s_n\sigma))$ with $\pi(s_i\sigma) \in \text{Val}_\pi$ for $i \in \{1, \dots, n\}$. Moreover suppose $\text{nrm}(f) = \{j_1, \dots, j_q\}$. Then*

$$f^n(N_\pi(s_{j_1}\sigma), \dots, N_\pi(s_{j_q}\sigma)) \succ_{3 \cdot \|\pi(t)\|} N_\pi(t\sigma)$$

holds.

Proof. Note that the assumption implies that the argument filtering π does not collapse f . We show the lemma by induction on $>_{\text{pop}}^\pi$. We consider the subcase that $s >_{\text{pop}}^\pi t$ follows as $t = g(t_1, \dots, t_m)$, π does not collapse on g , $f \succ g$, and $s >_{\text{pop}}^\pi t_j$ for all $j \in \pi(g)$, cf. Definition 4.1(ii). We set $u := f^n(N_\pi(s_{j_1}\sigma), \dots, N_\pi(s_{j_q}\sigma))$ and $k := 3 \cdot \|\pi(t)\|$ and first prove $u \succ_{k-1} S_\pi(t\sigma)$.

If $\pi(t\sigma) \in \text{Val}_\pi$, then $S_\pi(t\sigma) = []$ is minimal with respect to $\succ_{k-1}$. Thus we are done. Hence suppose $\text{nrm}(g) = \{j'_1, \dots, j'_q\}$, $\text{safe}(g) = \{i'_1, \dots, i'_p\}$ and let

$$S_\pi(t\sigma) = [g^n(N_\pi(t_{j'_1}\sigma), \dots, N_\pi(t_{j'_q}\sigma)) S_\pi(t_{i'_1}\sigma) \cdots S_\pi(t_{i'_p}\sigma)] .$$

We set $v := g^n(N_\pi(t_{j'_1}\sigma), \dots, N_\pi(t_{j'_q}\sigma))$. It suffices to show $u \succ_{k-2} v$ and $u \succ_{k-2} S_\pi(t_j\sigma)$ for $j \in \text{safe}(g)$. Both assertions follow from the induction hypothesis.

Now consider $N_\pi(t\sigma) = [S_\pi(t\sigma) \mathbf{s} \cdots \mathbf{s}]$ with $\|\pi(t\sigma)\|$ occurrences of the constant $\mathbf{s}$. Recall that $\text{width}(N_\pi(t\sigma)) = \|\pi(t\sigma)\| + 1$. Observe that $f^n \succ \mathbf{s}$. Hence to prove $u \succ_k S_\pi(t\sigma)$ it suffices to observe that $\text{width}(u) + k > \|\pi(t\sigma)\| + 1$ holds. For that note that $\|\pi(t\sigma)\|$ is either $\|\pi(t_j\sigma)\| + 1$ for some $j \in \pi(g)$ or less than k . In the latter case, we are done. Otherwise $\|\pi(t\sigma)\| = \|\pi(t_j\sigma)\| + 1$. Then from the definition of $\succ_k$ and the induction hypothesis $u \succ_{3 \cdot \|\pi(t_j)\|} N_\pi(t_j\sigma)$ we can conclude $\text{width}(u) + 3 \cdot \|\pi(t_j)\| > \text{width}(N_\pi(t_j\sigma)) = \|\pi(t_j\sigma)\| + 1$. Since $k \geq 3 \cdot (\|\pi(t_j)\| + 1)$, $\text{width}(u) + k > \|\pi(t\sigma)\| + 1$ follows. $\square$

Lemma 5.5. *Suppose $s >_{\text{pop}^*}^\pi t$ such that $\pi(s\sigma) = f(\pi(s_1\sigma), \dots, \pi(s_n\sigma))$ with $\pi(s_i\sigma) \in \text{Val}_\pi$ for $i \in \{1, \dots, n\}$. Then for $\text{nrm}(f) = \{j_1, \dots, j_q\}$,*

- (i) $f^n(\mathbf{N}_\pi(s_{j_1}\sigma), \dots, \mathbf{N}_\pi(s_{j_q}\sigma)) \blacktriangleright_{3 \cdot \|\pi(t)\|} \mathbf{S}_\pi(t\sigma)$, and
- (ii) $f^n(\mathbf{N}_\pi(s_{j_1}\sigma), \dots, \mathbf{N}_\pi(s_{j_q}\sigma)) :: \mathbf{BN}_\pi(s\sigma) \blacktriangleright_{3 \cdot \|\pi(t)\|} \mathbf{N}_\pi(t\sigma)$.

Proof. The lemma is shown by induction on the definition of $>_{\text{pop}^*}^\pi$. For the following, we set $u = f^n(\mathbf{N}_\pi(s_{j_1}\sigma), \dots, \mathbf{N}_\pi(s_{j_q}\sigma))$. Suppose $s >_{\text{pop}^*}^\pi t$ follows due to Definition 4.2(ii). We set $k := 3 \cdot \|\pi(t)\|$. Let $\text{nrm}(g) = \{j'_1, \dots, j'_q\}$ and let $\text{safe}(g) = \{i'_1, \dots, i'_p\}$.

Property (i) is immediate for $\pi(t\sigma) \in \text{Val}_\pi$, so assume otherwise. We see that $s >_{\text{pop}}^\pi t_j$ for all $j \in \text{nrm}(g)$ and obtain $u >_{k-1} g^n(\mathbf{N}_\pi(t_{j'_1}\sigma), \dots, \mathbf{N}_\pi(t_{j'_q}\sigma))$ as in Lemma 5.4. Furthermore, $s >_{\text{pop}^*}^\pi t_{j_0}$ for some $j_0 \in \text{safe}(g)$ and by induction hypothesis: $u \blacktriangleright_{k-1} \mathbf{S}_\pi(t_{j_0}\sigma)$. To conclude property (i), it remains to verify $u >_{k-1} \mathbf{S}_\pi(t_j\sigma)$ for the remaining $j \in \text{safe}(g)$. We either have $s >_{\text{pop}}^\pi t_j$ or $\pi(s_i) \supseteq \pi(t_j)$ (for some i). In the former subcase we proceed as in the claim, and for the latter we observe $\pi(t_j\sigma) \in \text{Val}_\pi$, and thus $\mathbf{S}_\pi(t_j\sigma) = []$ follows. This establishes property (i).

To conclude property (ii), it suffices to show $\text{width}(u :: \mathbf{BN}_\pi(s\sigma)) + k > \text{width}(\mathbf{N}_\pi(t\sigma))$, or equivalently $\|\pi(s\sigma)\| + 1 + k > \|\pi(t\sigma)\|$. The latter can be shown, if we proceed similar as in the claim. $\square$

Recall the definition of $\mathcal{Q}_{\mathcal{R}}$ from Section 2 and define $\mathcal{Q} := \{f(x_1, \dots, x_n) \rightarrow \perp \mid f \in \mathcal{D}\}$, and set $\mathcal{V}_{\mathcal{R}} := \mathcal{Q}_{\mathcal{R}}$. As the normal forms of $\mathcal{Q}$ coincide with Val , $\mathcal{V}_{\mathcal{R}}$ is the restriction of $\mathcal{I}_{\mathcal{R}}$, where arguments need to be values instead of normal forms of $\mathcal{R}$. From Lemma 5.3 and 5.5 we derive an embedding of root steps $\mathcal{V}_{\mathcal{R}}^\varepsilon$.

Suppose the step $s \mathcal{V}_{\mathcal{R}} t$ takes place below the root. Observe that $\pi(s) \neq \pi(t)$ need not hold in general. Thus we cannot hope to prove $\mathbf{N}_\pi(s) \blacktriangleright_k \mathbf{N}_\pi(t)$. However, we have the following stronger result.

Lemma 5.6. *There exists a uniform $k \in \mathbb{N}$ (depending only on $\mathcal{R}$) such that if $\mathcal{R} \subseteq >_{\text{pop}^*}^\pi$ holds then $s \mathcal{V}_{\mathcal{R}}^\varepsilon t$ implies $\mathbf{N}_\pi(s) \blacktriangleright_k \mathbf{N}_\pi(t)$. Moreover, if $\mathcal{R} \subseteq \gtrsim_{\text{pop}^*}^\pi$ holds then $s \mathcal{V}_{\mathcal{R}} t$ implies $\mathbf{N}_\pi(s) \blacktriangleright_k \mathbf{N}_\pi(t)$.*

Proof. We consider the first half of the assertion. Suppose $\mathcal{R} \subseteq >_{\text{pop}^*}^\pi$ and $s \mathcal{V}_{\mathcal{R}}^\varepsilon t$, that is for some rule $f(l_1, \dots, l_n) \rightarrow r \in \mathcal{R}$ and substitution $\sigma : \mathcal{V} \rightarrow \text{Val}$ we have $s = f(l_1\sigma, \dots, l_n\sigma)$ and $t = r\sigma$. Depending on whether π collapses f , the property either directly follows from Lemma 5.3 or is a consequence of Lemma 5.5(ii).

In order to conclude the second half of the assertion, one performs induction on the rewrite context. In addition, one shows that for the special case $\mathbf{S}_\pi(s) \approx \mathbf{S}_\pi(t)$, still $\|\pi(s)\| \geq \|\pi(t)\|$ holds. From this the lemma follows. $\square$

For constructor TRSs, we can simulate $\mathcal{I}_{\mathcal{R}}$ using $\mathcal{V}_{\mathcal{R}}$. We extend $\mathcal{R}$ with suitable rules $\Phi(\mathcal{R})$, which replace normal forms that are not values by some constructor symbol. To simplify the argument we re-use the symbol $\perp$ from above. We define the TRS $\Phi(\mathcal{R})$ as

$$\Phi(\mathcal{R}) := \{f(t_1, \dots, t_n) \rightarrow \perp \mid f(t_1, \dots, t_n) \in \text{NF}(\mathcal{R}) \cap \mathcal{T}(\mathcal{F}) \text{ and } f \in \mathcal{D}\}.$$

Moreover, we define $\phi_{\mathcal{R}}(t) := t \downarrow_{\Phi(\mathcal{R})}$. Observe that $\phi_{\mathcal{R}}(\cdot)$ is well-defined since $\Phi(\mathcal{R})$ is confluent and terminating.

Lemma 5.7. *Let $\mathcal{R} \cup \mathcal{S}$ be a constructor TRS. Define $\mathcal{S}' := \mathcal{S} \cup \Phi(\mathcal{R} \cup \mathcal{S})$. For $s \in \mathcal{T}(\mathcal{F})$,*

$$s \xrightarrow{\varepsilon}_{\mathcal{R}/\mathcal{S}} t \quad \text{implies} \quad \phi_{\mathcal{R} \cup \mathcal{S}}(s) \xrightarrow{\varepsilon}_{\mathcal{R}/\mathcal{S}'} \phi_{\mathcal{R} \cup \mathcal{S}}(t),$$

where $\xrightarrow{\varepsilon}_{\mathcal{R}/\mathcal{S}'}$ abbreviates $\xrightarrow{*}_{\mathcal{S}'} \cdot \xrightarrow{\varepsilon}_{\mathcal{R}} \cdot \xrightarrow{*}_{\mathcal{S}'}$.

Proof. It is easy to see that $s \xrightarrow{\varepsilon}_{\mathcal{R}} t$ implies $\phi_{\mathcal{R}}(s) \xrightarrow{\varepsilon}_{\mathcal{R}} \cdot \xrightarrow{\varepsilon}_{\Phi(\mathcal{R})} \phi_{\mathcal{R}}(t)$. Suppose $s \xrightarrow{\varepsilon}_{\mathcal{R}/\mathcal{S}} t$, then there exist ground terms u and v such that $s \xrightarrow{*}_{\mathcal{S}} u \xrightarrow{\varepsilon}_{\mathcal{R}} v \xrightarrow{*}_{\mathcal{S}} t$. Let $\phi(t) := \phi_{\mathcal{R} \cup \mathcal{S}}(t)$. From the above, $\phi(s) \xrightarrow{*}_{\mathcal{S}'} \phi(u) \xrightarrow{\varepsilon}_{\mathcal{R}} \cdot \xrightarrow{*}_{\mathcal{S}'} \phi(v) \xrightarrow{*}_{\mathcal{S}'} \phi(t)$ follows as desired. $\square$

Suppose $\mathcal{R} \subseteq >^{\pi}_{\text{pop}*}$ and $\mathcal{S} \subseteq \gtrsim^{\pi}_{\text{pop}*}$ holds. Together with Lemma 5.6, the above simulation establishes the promised embedding of $\xrightarrow{\varepsilon}_{\mathcal{R}/\mathcal{S}}$ into $\blacktriangleright_k$.

Lemma 5.8. *Let $\mathcal{R} \cup \mathcal{S}$ be a constructor TRS, and suppose $\mathcal{R} \subseteq >^{\pi}_{\text{pop}*}$ and $\mathcal{S} \subseteq \gtrsim^{\pi}_{\text{pop}*}$ hold. Then for k depending only on $\mathcal{R}$ and $\mathcal{S}$ and $s \in \mathcal{T}(\mathcal{F})$, we have*

$$s \xrightarrow{\varepsilon}_{\mathcal{R}/\mathcal{S}} t \quad \text{implies} \quad \mathbf{N}_{\pi}(\phi(s)) \blacktriangleright_k^+ \mathbf{N}_{\pi}(\phi(t)).$$

Proof. Consider a step $s \xrightarrow{\varepsilon}_{\mathcal{R}/\mathcal{S}} t$ and set $\phi(t) := \phi_{\mathcal{R} \cup \mathcal{S}}(t)$. By Lemma 5.7 there exist terms u and v such that $\phi(s) \xrightarrow{*}_{\mathcal{S} \cup \Phi(\mathcal{R} \cup \mathcal{S})} u \xrightarrow{\varepsilon}_{\mathcal{R}} v \xrightarrow{*}_{\mathcal{S} \cup \Phi(\mathcal{R} \cup \mathcal{S})} \phi(t)$. Since $\mathcal{R} \subseteq >^{\pi}_{\text{pop}*}$ holds, by Lemma 5.6 $\mathbf{N}_{\pi}(u) \blacktriangleright_{k_1} \mathbf{N}_{\pi}(v)$ follows. Moreover from $\mathcal{S} \subseteq \gtrsim^{\pi}_{\text{pop}*}$ together with Lemma 5.6 we conclude that $r_1 \xrightarrow{\varepsilon}_{\mathcal{S} \cup \Phi(\mathcal{R} \cup \mathcal{S})} r_2$ implies $\mathbf{N}_{\pi}(r_1) \blacktriangleright_{k_2} \mathbf{N}_{\pi}(r_2)$. Here it suffices to see that steps from $\mathcal{V}(\mathcal{R} \cup \mathcal{S})$ are easy to embed into $\blacktriangleright_{k_2}$ using the predicative interpretation $\mathbf{N}_{\pi}$ independent of k_2 . In both cases k_1 and k_2 depend only on $\mathcal{R}$ and $\mathcal{S}$ respectively; set $k := \max\{k_1, k_2\}$. In sum we have $\mathbf{N}_{\pi}(\phi(s)) \blacktriangleright_k^* \mathbf{N}_{\pi}(u) \blacktriangleright_k \mathbf{N}_{\pi}(v) \blacktriangleright_k^* \mathbf{N}_{\pi}(\phi(t))$, employing $\blacktriangleright_{l_1} \subseteq \blacktriangleright_{l_2}$ for $l_1 \leq l_2$. It is an easy exercise to show that $\blacktriangleright_k \cdot \approx \subseteq \blacktriangleright_k$ and likewise $\approx \cdot \blacktriangleright_k \subseteq \blacktriangleright_k$ holds. Hence the lemma follows. $\square$

Theorem 5.9. *Let $\mathcal{R} \cup \mathcal{S}$ be a constructor TRS, and suppose $\mathcal{R} \subseteq >^{\pi}_{\text{pop}*}$ and $\mathcal{S} \subseteq \gtrsim^{\pi}_{\text{pop}*}$ holds. Then there exists a polynomial p depending only on $\mathcal{R} \cup \mathcal{S}$ such that for any basic and ground term t , $\text{dl}(t, \xrightarrow{\varepsilon}_{\mathcal{R}/\mathcal{S}}) \leq p(|t|)$.*

Proof. Assume $t \notin \text{NF}(\mathcal{R} \cup \mathcal{S})$, otherwise $\text{dl}(t, \xrightarrow{\varepsilon}_{\mathcal{R}/\mathcal{S}})$ is trivially bounded. Moreover t is a basic term, hence $\phi_{\mathcal{R} \cup \mathcal{S}}(t) = t$. From Lemma 5.8 we infer that $\text{dl}(t, \xrightarrow{\varepsilon}_{\mathcal{R}/\mathcal{S}}) \leq \mathbf{G}_k(\mathbf{N}_{\pi}(\phi_{\mathcal{R} \cup \mathcal{S}}(t))) = \mathbf{G}_k(\mathbf{N}_{\pi}(t))$ for some k , where the latter is polynomially bounded in $|t|$ and the polynomial only depends on k , cf. Lemma 5.2. Finally k depends only on $\mathcal{R} \cup \mathcal{S}$. $\square$

Suppose $\mathcal{R}$ is a constructor TRS, and let $\mathcal{P}$ denote the set of weak innermost dependency pairs. For the moment, suppose that all compound symbols of $\mathcal{P}$ are nullary. Provided that $\mathcal{P}$ is non-duplicating and compatible with some SLI, as a consequence of the above theorem paired with Proposition 3.4, the inclusions $\mathcal{P} \subseteq >^{\pi}_{\text{pop}*}$ and $\mathcal{U}(\mathcal{P}) \subseteq \gtrsim^{\pi}_{\text{pop}*}$ certify that $\text{rc}_{\mathcal{R}}^{\dagger}$ is polynomially bounded. Observe that for the application of $>^{\pi}_{\text{pop}*}$ and $\gtrsim^{\pi}_{\text{pop}*}$ in the context of $\mathcal{P}$ and $\mathcal{U}(\mathcal{P})$, we alter Definitions 4.1 and 4.2 such that $f \in \mathcal{D}^{\sharp}$ is demanded.

Example 5.10 (Example 4.5 continued). Reconsider the TRS $\mathcal{R}_{\text{bits}}$, and let $\mathcal{P}$ denote $\text{WIDP}(\mathcal{R}_{\text{bits}})$ as drawn in Example 3.2. By taking the SLI $\mathcal{A}$ with $0_{\mathcal{A}} = 0$, $s_{\mathcal{A}}(x) = x + 1$ and $\text{half}_{\mathcal{A}}(x) = x + 1$ we obtain $\mathcal{U}(\mathcal{P}) \subseteq >_{\mathcal{A}}$ and moreover, observe that $\mathcal{P}$ is both non-duplicating and contains only nullary compound symbols. In Example 4.5 we have seen that $\mathcal{P} \subseteq >_{\text{pop}^*}^{\pi}$ holds. Similar, $\mathcal{U}(\text{WIDP}(\mathcal{R}_{\text{bits}})) \subseteq \gtrsim_{\text{pop}^*}^{\pi}$ can easily be shown. From the above observation we thus conclude a polynomial runtime-complexity of $\mathcal{R}_{\text{bits}}$.

The assumption that all compound symbols from $\mathcal{P}$ need to be nullary is straightforward to lift, but technical. Hence, we do not provide a complete proof here, but only indicate the necessary changes. The formal construction can be found in the Appendix.

Note that in the general case, it does not suffice to embed root steps of $\mathcal{P}$ into $\blacktriangleright_k$, rather we have to embed steps of form $C[s_1^{\sharp}, \dots, s_i^{\sharp}, \dots, s_n^{\sharp}] \xrightarrow{\mathcal{P}} C[s_1^{\sharp}, \dots, t_i^{\sharp}, \dots, s_n^{\sharp}]$ with C being a context built from compound symbols. As first measure we require that the argument filtering π is *safe* [15], that is $\pi(c) = [1, \dots, n]$ for each compound symbol c of arity n . Secondly, we adapt the predicative interpretation $\mathbf{N}_{\pi}$ in such a way that compound symbols are interpreted as sequences, and their arguments by the interpretation $\mathbf{N}_{\pi}$. This way, a proper embedding using $\mathbf{N}_{\pi}$ requires $\mathbf{N}_{\pi}(s_i^{\sharp}) \blacktriangleright_k \mathbf{N}_{\pi}(t_i^{\sharp})$ instead of $\mathbf{S}_{\pi}(s_i^{\sharp}) \blacktriangleright_k \mathbf{S}_{\pi}(t_i^{\sharp})$.

Theorem 5.11. *Let $\mathcal{R}$ be a constructor TRS, and let $\mathcal{P}$ denote the set of weak innermost dependency pairs. Assume $\mathcal{P}$ is non-duplicating, and suppose $\mathcal{U}(\mathcal{P}) \subseteq >_{\mathcal{A}}$ for some SLI $\mathcal{A}$. Let π be a safe argument filtering. If $\mathcal{P} \subseteq >_{\text{pop}^*}^{\pi}$ and $\mathcal{U}(\mathcal{P}) \subseteq \gtrsim_{\text{pop}^*}^{\pi}$ then $\text{rc}_{\mathcal{R}}^{\dagger}$ is polynomially bounded.*

Above it is essential that $\mathcal{R}$ is a constructor TRS. This even holds when POP^* is applied directly.

Example 5.12. Consider the TRS $\mathcal{R}_{\text{exp}}$ below:

$$\begin{aligned} \text{exp}(x) &\rightarrow \text{e}(\text{g}(x)) & \text{e}(\text{g}(s(x))) &\rightarrow \text{dp}_1(\text{g}(x)) & \text{g}(0) &\rightarrow 0 \\ \text{dp}_1(x) &\rightarrow \text{dp}_2(\text{e}(x), x) & \text{dp}_2(x, y) &\rightarrow \text{pr}(x, \text{e}(y)) \end{aligned}$$

The above rules are oriented (directly) by $>_{\text{pop}^*}$ induced by **safe** and $\gtrsim$ such that: (i) the argument position of **g** and **exp** are normal, the remaining argument positions are safe, and (ii) $\text{exp} \succ \text{g} \succ \text{dp}_1 \succ \text{dp}_2 \succ \text{e} \succ \text{pr} \succ 0$. On the other hand, $\mathcal{R}_{\text{exp}}$ admits at least exponential innermost runtime-complexity, as for instance $\text{exp}(s^n(0))$ normalizes in exponentially (in n) many innermost rewrite steps.

To overcome this obstacle, we adapt the definition of $>_{\text{pop}^*}$ in the sense that we refine the notion of defined function symbols as follows. Let $\mathcal{G}_{\mathcal{C}}$ denote the least set containing $\mathcal{C}$ and all symbols appearing in arguments to left-hand sides in $\mathcal{R}$. Moreover, set $\mathcal{G}_{\mathcal{D}} := \mathcal{F} \setminus \mathcal{G}_{\mathcal{C}}$ and set $\text{Val} := \mathcal{T}(\mathcal{G}_{\mathcal{C}}, \mathcal{V})$. Then in order to extend Theorem 5.11 to non-constructor TRS it suffices to replace $\mathcal{D}$ by $\mathcal{G}_{\mathcal{D}}$ and $\mathcal{C}$ by $\mathcal{G}_{\mathcal{C}}$ in all above given definitions and arguments (see [3] for the formal construction). Thus the next theorem follows easily from combining Proposition 3.6 and Theorem 5.11. Note that this theorem can be easily extended so that in each path different termination techniques (inducing polynomial runtime complexity) are employed, see [14] and Section 6.

Theorem 5.13. *Let $\mathcal{R}$ be a TRS. Let $\mathcal{G}$ denote the weak innermost dependency graph, and let $\mathcal{F} = \mathcal{G}_D \uplus \mathcal{G}_C$ be separated as above. Suppose for every path $(\mathcal{P}_1, \dots, \mathcal{P}_n)$ in $\mathcal{G}/\equiv$ there exists an SLI $\mathcal{A}$ and a pair $(\succsim_{\text{pop}^*}^\pi, >_{\text{pop}^*}^\pi)$ based on a safe argument filtering π such that (i) $\mathcal{U}(\mathcal{P}_1 \cup \dots \cup \mathcal{P}_n) \subseteq >_{\mathcal{A}}$ (ii) $\mathcal{P}_1 \cup \dots \cup \mathcal{P}_{n-1} \cup \mathcal{U}(\mathcal{P}_1 \cup \dots \cup \mathcal{P}_n) \subseteq \succsim_{\text{pop}^*}^\pi$, and (iii) $\mathcal{P}_n \subseteq >_{\text{pop}^*}^\pi$ holds. Then $\text{rc}_{\mathcal{R}}^i$ is polynomially bounded.*

The next theorem establishes that POP* in conjunction with (weak innermost) dependency pairs induces polytime computability of the function described through the analysed TRS. We kindly refer the reader to the Appendix for the proof.

Theorem 5.14. *Let $\mathcal{R}$ be an orthogonal, S -sorted and completely defined constructor TRS such that the underlying signature is simple. Let $\mathcal{P}$ denote the set of weak innermost dependency pairs. Assume $\mathcal{P}$ is non-duplicating, and suppose $\mathcal{U}(\mathcal{P}) \subseteq >_{\mathcal{A}}$ for some SLI $\mathcal{A}$. If $\mathcal{P} \subseteq \succsim_{\text{pop}^*}^\pi$ and $\mathcal{U}(\mathcal{P}) \subseteq \succsim_{\text{pop}^*}^\pi$ then the functions computed by $\mathcal{R}$ are computable in polynomial time.*

Here *simple* signature [18] essentially means that the size of any constructor term depends polynomially on its depth. Such a restriction is always necessary in this context. A detailed account is given in the Appendix (see also [18]). This restriction is also responsible for the introduction of sorts.

6. Experimental Results

All described techniques have been incorporated into the *Tyrolean Complexity Tool* TCT, an open source complexity analyser². We performed tests on two testbeds: **T** constitutes of the 1394 examples from the Termination Problem Database Version 5.0.2 that were used in the runtime-complexity category of the termination competition 2008³. Moreover, testbed **C** is the restriction of testbed **T** to constructor TRSs (638 in total). All experiments were conducted on a machine that is identical to the official competition server (8 AMD Opteron[®] 885 dual-core processors with 2.8GHz, 8x8 GB memory). As timeout we use 5 seconds. We orient TRSs using $>_{\text{pop}^*}^\pi$ by encoding the constraints on precedence and so forth in *propositional logic* (cf. [3] for details), employing MiniSat [9] for finding satisfying assignments. In a similar spirit, we check compatibility with SLIs via translations to SAT. In order to derive an estimated dependency graph, we use the function ICAP (cf. [13]).

Experimental findings are summarised in Table 1.⁴ In each column, we highlight the total on yes-, maybe- and timeout-instances. Furthermore, we annotate average times in seconds. In the first three columns we contrast POP* as direct technique to POP* as base to (weak innermost) dependency pairs. I.e., the columns WIDP and WIDG show results concerning Proposition 3.4 together with Theorem 5.11 or Theorem 5.13 respectively.

In the remaining four columns we assess the power of Proposition 3.4 and 3.6 in conjunction with different base orders, thus verifying that the use of POP* in this context

² Available at <http://cl-informatik.uibk.ac.at/software/tct>.

³ See <http://termcomp.uibk.ac.at>.

⁴ See <http://cl-informatik.uibk.ac.at/~zini/rta09> for extended results.

is independent to existing techniques. Column P asserts that the different paths are handled by *linear and quadratic restricted interpretations* [15]. In column PP, in addition POP* is employed. Similar, in column M *restricted matrix interpretations* (that is matrix interpretations [10], where constructors are interpreted by triangular matrices) are used to handle different paths. Again column MP extends column M with POP*. Note that all methods induce polynomial innermost runtime complexity.

		polynomial path orders			dependency graphs mixed			
		DIRECT	WIDP	WIDG	P	PP	M	MP
T	Yes	46/0.03	69/0.09	80/0.07	198/0.54	198/0.51	200/0.63	207/0.48
	Maybe	1348/0.04	1322/0.10	1302/0.14	167/0.77	170/0.82	142/0.61	142/0.63
	Timeout	0	3	12	1029	1026	1052	1045
C	Yes	40/0.03	48/0.08	55/0.05	99/0.40	100/0.38	98/0.26	105/0.23
	Maybe	598/0.05	587/0.10	576/0.13	143/0.72	146/0.77	119/0.51	119/0.54
	Timeout	0	3	7	396	392	421	414

Table 1: Experimental Results

Table 1 reflects that the integration of POP* in the context of (weak) dependency pairs, significantly extends the direct approach. Worthy of note, the extension of [4] with quasi-precedences alone gives 5 additional examples. As advertised, POP* is incredibly fast in all settings. Consequently, as evident from the table, polynomial path orders team well with existing techniques, without affecting overall performance: notice that due to the additional of POP* the number of timeouts is reduced.

7. Conclusion

In this paper we study the runtime complexity of rewrite systems. We combine two recently developed techniques in the context of complexity analysis: weak innermost dependency pairs and polynomial path orders. If the conditions of our main result are met, we can conclude the innermost polynomial runtime complexity of the studied term rewrite system. And we obtain that the function defined are *polytime computable*. We have implemented the technique and experimental evidence clearly indicates the power and in particular the efficiency of the new method.

A. Appendix

Below we present the missing proofs of Theorem 5.11 and Theorem 5.14 respectively.

As mentioned in Section 5, we now introduce an *extended predicative interpretation* whose purpose is to interpret compound symbols as sequences, and their arguments via the interpretation N_π .

Definition A.1. The *extended predicative interpretation* $\mathbf{N}_\pi^s$ from terms $\mathcal{T}(\mathcal{F}, \mathcal{V})$ to sequences $\text{Seq}(\mathcal{F}_\pi^n \cup \{\mathbf{s}\}, \mathcal{V})$ is defined as follows: if $t = \mathbf{c}(t_1, \dots, t_n)$ and $\mathbf{c} \in \mathcal{C}_{com}$ then $\mathbf{N}_\pi^s(t) := [\mathbf{N}_\pi^s(t_1) \ \dots \ \mathbf{N}_\pi^s(t_n)]$, and otherwise $\mathbf{N}_\pi^s(t) := [\mathbf{N}_\pi(t)]$.

Following [20, Section 6.5], we briefly recall *typed rewriting*. Let S be a finite set representing the set of *types* or *sorts*. An *S-sorted set* A is a family of sets $\{A_s \mid s \in S\}$ such that all sets A_s are pairwise disjoint. In the following, we suppose that $\mathcal{V}$ denotes an *S-sorted set* of variables. An *S-sorted signature* $\mathcal{F}$ is like a signature, but the *arity* of $f \in \mathcal{F}$ is defined by $\text{ar}(f) = (s_1, \dots, s_n)$ for $s_1, \dots, s_n \in S$. Additionally, each symbol $f \in \mathcal{F}$ is associated with a sort $s \in S$, called the *type of f* and denoted by $\text{st}(f)$. We adopt the usual notion and write $f : (s_1, \dots, s_n) \rightarrow s$ when $\text{ar}(f) = (s_1, \dots, s_n)$ and $\text{st}(f) = s$. The *S-sorted set of terms* $\mathcal{T}(\mathcal{F}, \mathcal{V})_S$ consists of the sets $\mathcal{T}(\mathcal{F}, \mathcal{V})_s$ for $s \in S$, where $\mathcal{T}(\mathcal{F}, \mathcal{V})_s$ is inductively defined by (i) $\mathcal{V}_s \subseteq \mathcal{T}(\mathcal{F}, \mathcal{V})_s$, and (ii) $f(t_1, \dots, t_n) \in \mathcal{T}(\mathcal{F}, \mathcal{V})_s$ for all function symbols $f \in \mathcal{F}$, $f : (s_1, \dots, s_n) \rightarrow s$ and terms $t_i \in \mathcal{T}(\mathcal{F}, \mathcal{V})_{s_i}$ for $i \in \{1, \dots, n\}$. We say that a term t is *well-typed* if $t \in \mathcal{T}(\mathcal{F}, \mathcal{V})_s$ for some sort s . An *S-sorted term rewrite system* $\mathcal{R}$ is a TRS such that for $l \rightarrow r \in \mathcal{R}$, it holds that $l, r \in \mathcal{T}(\mathcal{F}, \mathcal{V})_s$ for some sort $s \in S$. As a consequence, for $s \in \mathcal{T}(\mathcal{F}, \mathcal{V})_s$ and $s \rightarrow_{\mathcal{R}} t$, we have that $t \in \mathcal{T}(\mathcal{F}, \mathcal{V})_s$.

Example A.2. Let $S = \{\text{Bool}, \text{List}, \text{Nat}, \text{Pair}\}$. The *S-sorted* rewrite system $\mathcal{R}_{\text{Lst}}$ is given by the following rules:

$$\begin{array}{ll} \mathbf{f}(\mathbf{s}(x)) \rightarrow \text{cons}(\text{pair}(x, \mathbf{g}(x)), \mathbf{f}(x)) & \mathbf{g}(\mathbf{s}(x)) \rightarrow \mathbf{g}(x) \\ \mathbf{f}(0) \rightarrow \text{nil} & \mathbf{g}(0) \rightarrow \text{tt} \end{array}$$

Here we assign arities and sorts as follows: for the constructors we set $0 : \text{Nat}$, $\mathbf{s} : \text{Nat} \rightarrow \text{Nat}$, $\text{pair} : (\text{Nat}, \text{Bool}) \rightarrow \text{Pair}$, $\text{tt} : \text{Bool}$, $\text{nil} : \text{List}$, $\text{cons} : (\text{Pair}, \text{List}) \rightarrow \text{List}$; for the defined symbols we set $\mathbf{f} : \text{Nat} \rightarrow \text{List}$ and $\mathbf{g} : \text{Nat} \rightarrow \text{Bool}$.

A *simple* signature [18] is a sorted signature such that each sort has a finite *rank* r in the following sense: the sort s has rank r if for every constructor $c : (s_1, \dots, s_n) \rightarrow s$, the rank of each sort s_i is less than the rank of s , except for at most one sort which can be of rank r . Simple signatures allow the definition of enumerated datatypes and inductive datatypes like words and lists but prohibit for instance the definition of tree structures. Observe that the signature underlying $\mathcal{R}_{\text{Lst}}$ from Example A.2 is simple. A crucial insight is that sizes of values formed from a simple signature can be estimated polynomially in their depth. The easy proof of the following proposition can be found in [18, Proposition 17].

Proposition A.3. *Let $\mathcal{C}$ be a set of constructors from a simple signature $\mathcal{F}$. There exists a constant $d \in \mathbb{N}$ such that for each term $t \in \mathcal{T}(\mathcal{C}, \mathcal{V})_S$ whose rank is r , $|t| \leq d^r \cdot \text{dp}(t)^{r+1}$.*

In order to give a polytime algorithm for the functions computed by a TRS, it is essential that sizes of reducts do not exceed a polynomial bound with respect to the size of the start term. Recall that approximations $\blacktriangleright_k$ tightly control the size growth of terms. For simple signatures, we can exploit this property for a space-complexity

analysis. Although predicative interpretations remove values, by the above proposition sizes of those can be estimated based on the Buchholz-norm record in $\mathbf{N}_\pi$. And so we derive the following Lemma, essential for the proof of Theorem 5.14.

Lemma A.4. *Let $\mathcal{F}$ be a simple signature. There exists a (monotone) polynomial p depending only on $\mathcal{F}$ such that for each well-typed term $t \in \mathcal{T}(\mathcal{F}, \mathcal{V})_s$, $|t| \leq p(\mathbf{G}_k(\mathbf{N}^s(t)))$.*

Proof. The Lemma follows as: (i) for all sequences $s \in \mathcal{Seq}$, $|s| \leq \mathbf{G}_k(s) + 1$, and (ii) for all terms $t \in \mathcal{T}(\mathcal{F}, \mathcal{V})_s$, $|t| \leq c \cdot |\mathbf{N}^s(t)|^d$ for some uniform constants $0 < c, d \in \mathbb{N}$. These properties are simple to verify: property (i) follows from induction on s where we employ for the inductive step that $f(s_1, \dots, s_n) \blacktriangleright_k [s_1 \cdots s_n]$ and $\mathbf{G}_k([s_1 \cdots s_n]) = \sum_{i=1}^n \mathbf{G}_k(s_i) + n$. For property (ii), set $d = r + 2$ where r is the maximal rank of a symbol in $\mathcal{C}$, and set $c = e^r$ where e is as given from Proposition A.3. First one shows by a straight forward induction on t that $|t| \leq c \cdot (|\mathbf{S}(t)| \cdot \|t\|^{r+1})$ (employing Proposition A.3 and $\mathbf{dp}(t) \leq \|t\|$). As $|\mathbf{S}(t)| < |\mathbf{N}(t)|$ and $\|t\| < |\mathbf{N}(t)|$, we derive $|t| < c \cdot |\mathbf{N}(t)|^d$. By induction on the definition of $\mathbf{N}^s$ we finally obtain property (ii). $\square$

Let $\mathcal{R}$ be a (not necessarily S -sorted) TRS that is innermost terminating. In the sequel, we keep $\mathcal{R}$ fixed. In order to exploit Lemma A.4 for an analysis by means of weak innermost dependency pairs, we introduce the notion of *type preserving weak innermost dependency pairs*.

Definition A.5. If $l \rightarrow r \in \mathcal{R}$ and $r = C\langle u_1, \dots, u_n \rangle_{\mathcal{D}}$ then $l^\# \rightarrow \mathbf{c}(u_1^\#, \dots, u_n^\#)$ is called a *type preserving weak innermost dependency pair* of $\mathcal{R}$. Here, the *compound symbol* $\mathbf{c}$ is supposed to be fresh. We set $\mathbf{repr}(\mathbf{c}) := C$ and say that $\mathbf{c}$ *represents* the context C . The set of all type preserving weak innermost dependency pairs is denoted by $\mathbf{WIDP}(\mathcal{R})$.

We collect all compound symbols appearing in $\mathbf{TPWIDP}(\mathcal{R})$ in the set $\mathcal{C}_{com}$.

Example A.6 (Example A.2 continued). Reconsider the rewrite system $\mathcal{R}_{\text{Lst}}$ given in Example A.2. The set $\mathbf{TPWIDP}(\mathcal{R}_{\text{Lst}})$ is given by

$$\begin{aligned} f^\#(s(x)) &\rightarrow \mathbf{c}_1(g^\#(x), f^\#(x)) & g^\#(s(x)) &\rightarrow \mathbf{c}_3(g^\#(x)) \\ f^\#(0) &\rightarrow \mathbf{c}_2 & g^\#(0) &\rightarrow \mathbf{c}_4 \end{aligned}$$

The constant $\mathbf{c}_3$ represents for instance the empty context, and the constant $\mathbf{c}_1$ represents the context $\mathbf{repr}(\mathbf{c}_1) = \mathbf{cons}(\mathbf{pair}(x, \square), \square)$.

Lemma A.7. *Let $\mathcal{R}$ be an S -sorted TRS such that the underlying signature $\mathcal{F}$ is simple. Then $\mathbf{TPWIDP}(\mathcal{R}) \cup \mathcal{U}(\mathbf{WIDP}(\mathcal{R}))$ is an S -sorted TRS, and the underlying signature $\mathcal{F}^\# \cup \mathcal{C}_{com}$ a simple signature.*

Proof. To conclude the claim, it suffices to type the marked and compound symbols appropriately. For each rule $f^\#(l_1, \dots, l_n) \rightarrow \mathbf{c}(r_1^\#, \dots, r_n^\#) \in \mathbf{TPWIDP}(\mathcal{R})$ we proceed as follows: we set $\mathbf{ar}(f^\#) := \mathbf{ar}(f)$ and $\mathbf{st}(f^\#) := \mathbf{st}(f)$. Moreover, we set $\mathbf{ar}(\mathbf{c}) := (\mathbf{st}(r_1), \dots, \mathbf{st}(r_m))$ and $\mathbf{st}(\mathbf{c}) := \mathbf{st}(f)$. It is easy to see that since $\mathcal{R}$ is S -sorted, $\mathbf{TPWIDP}(\mathcal{R}) \cup \mathcal{U}(\mathbf{TPWIDP}(\mathcal{R}))$ is S -sorted too. $\square$

Note that the above lemma fails for weak innermost dependency pairs: consider the rule $f(x) \rightarrow d(g(x))$, where f and g are defined symbols and d is a constructor. Moreover, suppose $f : s_2 \rightarrow s_1$, $g : s_2 \rightarrow s_3$ and $d : s_3 \rightarrow s_1$. Then we cannot type the corresponding weak innermost dependency pair $f^\sharp(x) \rightarrow g^\sharp(x)$ as above because (return-)types of $f^\sharp$ and $g^\sharp$ differ.

As for practical all termination techniques, compatibility of weak innermost dependency pairs with polynomial path orders also yield compatibility of type preserving weak innermost dependency pairs. Moreover, from the definition we immediately see that $dl(t^\sharp, \xrightarrow{\text{TPWIDP}(\mathcal{R})/\mathcal{U}}) = dl(t^\sharp, \xrightarrow{\text{WIDP}(\mathcal{R})/\mathcal{U}})$ with $\mathcal{U} = \mathcal{U}(\text{WIDP}(\mathcal{R}))$ and basic term t . And so it is clear that in order to proof Theorem 5.11 and Theorem 5.14, $\text{WIDP}(\mathcal{R})$ can safely be replaced by $\text{TPWIDP}(\mathcal{R})$. We continue with the proof of Theorem 5.11.

A.1. Proof of Theorem 5.11

Let ComCtx abbreviate the set of contexts $\mathcal{T}(\mathcal{C}_{\text{com}} \cup \{\square\}, \mathcal{V})$ build from compound symbols. Set $\mathcal{P} = \text{TPWIDP}(\mathcal{R})$ and $\mathcal{U} = \mathcal{U}(\text{WIDP}(\mathcal{R}))$. In order to highlight the correspondence between $\xrightarrow{\mathcal{R}}$ and $\xrightarrow{\mathcal{P}/\mathcal{U}}$, we extend the notion of *representatives*.

Definition A.8. Let $C \in \text{ComCtx}$. We define $\text{reprs}(C)$ as the least set of (ground) contexts such that (i) if $C = \square$ then $\square \in \text{reprs}(C)$, and (ii) if $C = c(C_1, \dots, C_n)$, $C'_i \in \text{reprs}(C_i)$ and σ is a substitution from all variables in $\text{repr}(c)$ to ground normal forms of $\mathcal{R}$ then $(\text{repr}(c)\sigma)[C'_1, \dots, C'_n] \in \text{reprs}(C)$.

Example A.9 (Example A.6 continued). Reconsider the TRS $\mathcal{R}_{\text{Lst}}$ from Example A.2, together with $\text{TPWIDP}(\mathcal{R}_{\text{Lst}})$ as given in Example A.6. Consider the step $f(s(0)) \rightarrow_{\mathcal{R}_{\text{Lst}}} \text{cons}(\text{pair}(0, g(0)), f(0))$ and the corresponding dependency pair step

$$f^\sharp(s(0)) \rightarrow_{\text{TPWIDP}(\mathcal{R}_{\text{Lst}})} c_1(g^\sharp(0), f^\sharp(0)) .$$

Let $C = c_1(\square, \square)$, remember that $\text{repr}(c_1) = \text{cons}(\text{pair}(x, \square), \square)$, $\text{reprs}(\square) = \square$ and observe that $C' = \text{cons}(\text{pair}(0, \square), \square) \in \text{reprs}(C)$ by taking the substitution $\sigma = \{x \mapsto 0\}$. And hence we can reformulate the above two steps as $f(s(0)) \rightarrow_{\mathcal{R}_{\text{Lst}}} C'[g(0), f(0)]$ and likewise $f^\sharp(s(0)) \rightarrow_{\text{TPWIDP}(\mathcal{R}_{\text{Lst}})} C[g^\sharp(0), f^\sharp(0)]$.

We manifest the above observation in the following lemma.

Lemma A.10. Let $s \in \mathcal{T}_{\text{b}}$ be a ground and basic term. Suppose $s \xrightarrow{\mathcal{R}}^* t$. Let $\mathcal{P} = \text{TPWIDP}(\mathcal{R})$ and let $\mathcal{U} = \mathcal{U}(\text{WIDP}(\mathcal{R}))$. Then there exists contexts $C' \in \text{ComCtx}$, $C \in \text{reprs}(C')$ and terms $t_1, \dots, t_n$ such that $t = C[t_1, \dots, t_n]$ and moreover, $s^\sharp \xrightarrow{\mathcal{P} \cup \mathcal{U}}^* C'[t_1^\sharp, \dots, t_n^\sharp]$.

Proof. We proof the lemma by induction on the length of the rewrite sequence $s \xrightarrow{\mathcal{R}}^n t$. The base case $n = 0$ is trivial, we set $C = C' = \square$. So suppose $s \xrightarrow{\mathcal{R}}^n t \xrightarrow{\mathcal{R}} u$ and the property holds for n . And thus we can find contexts $C'_t \in \text{ComCtx}$, $C_t \in \text{reprs}(C'_t)$ and terms $t_1, \dots, t_n$ such that $t = C_t[t_1, \dots, t_n]$ and moreover, $s^\sharp \xrightarrow{\mathcal{P} \cup \mathcal{U}}^* C'_t[t_1^\sharp, \dots, t_n^\sharp]$.

Without loss of generality we can assume $u = C_t[t_1, \dots, u_i, \dots, t_n]$ with $t_i \xrightarrow{i}_{\mathcal{R}} u_i$, as the context C_t is solely build from constructors and normal forms of $\mathcal{R}$.

First, suppose $t_i \xrightarrow{i}_{\mathcal{R}} u_i$, and hence $t_i = l\sigma$ for $l \rightarrow r \in \mathcal{R}$ and substitution $\sigma : \mathcal{V} \rightarrow \text{NF}(\mathcal{R}) \cap \mathcal{T}(\mathcal{F})$. Moreover $l^\# \rightarrow c(r_1^\#, \dots, r_m^\#) \in \mathcal{P}$ such that $u_i = (\text{repr}(c)\sigma)[r_1\sigma, \dots, r_m\sigma]$. We set C' as the context obtained from replacing the i -th hole of C_t' by $c(\square, \dots, \square)$, likewise we set C as the context obtained from replacing the i -th hole of C_t by $\text{repr}(c)\sigma$. Note that $C \in \text{reprs}(C')$. We conclude $s^\# \xrightarrow{i}_{\mathcal{P} \cup \mathcal{U}}^* C'[t_1^\#, \dots, r_1^\#\sigma, \dots, r_m^\#\sigma, \dots, t_n^\#]$ and $u = C[t_1, \dots, r_1\sigma, \dots, r_m\sigma, \dots, t_n]$ which establishes the lemma for this case.

Now suppose $t_i \xrightarrow{i}_{\mathcal{R}} u_i$ is a step below the root. Thus we have also $t_i^\# \xrightarrow{i}_{\mathcal{R}} u_i^\#$. As shown in [15, Lemma 16], the latter can be strengthened to $t_i^\# \xrightarrow{\mathcal{P} \cup \mathcal{U}}_{\mathcal{U}} u_i^\#$. We conclude $s^\# \xrightarrow{i}_{\mathcal{P} \cup \mathcal{U}}^* C_t'[t_1^\#, \dots, u_i^\#, \dots, t_n^\#]$, and the lemma follows by setting $C' = C_t'$ and $C = C_t$. $\square$

Suppose $\text{WIDP}(\mathcal{R})$ contains non-nullary compound symbols. In order to establish an embedding in the sense of Lemma 5.6 for that case, by the above lemma we see that it suffices to consider only terms of shape $s = C[s_1^\#, \dots, s_n^\#]$ with $C \in \text{ComCtx}$. With this insight, we adjust Lemma 5.6 as below. Observe that due to the definition of $\mathbf{N}_\pi^s$, we cannot simply apply Lemma 5.6 together with closure under context of $\blacktriangleright_k$ here.

Lemma A.11. *Let $s = C[s_1^\#, \dots, s_n^\#]$ for $C \in \text{ComCtx}$ and $s_1, \dots, s_n \in \mathcal{T}(\mathcal{F}, \mathcal{V})$. Let $\mathcal{P} = \text{TPWIDP}(\mathcal{R})$ and $\mathcal{U} = \mathcal{U}(\text{WIDP}(\mathcal{R}))$. There exists a uniform constant $k \in \mathbb{N}$ depending only on $\mathcal{R}$ such that if $\mathcal{P} \subseteq >_{\text{pop}^*}^\pi$ holds then $s \xrightarrow{\mathcal{P}} t$ implies $\mathbf{N}_\pi^s(s) \blacktriangleright_k \mathbf{N}_\pi^s(t)$. Moreover, if $\mathcal{U} \subseteq \gtrsim_{\text{pop}^*}^\pi$ holds then $s \xrightarrow{\mathcal{U}} t$ implies $\mathbf{N}_\pi^s(s) \blacktriangleright_k \mathbf{N}_\pi^s(t)$.*

Proof. We proof the lemma for $k := \max\{3 \cdot \|r\| \mid l \rightarrow r \in \mathcal{P} \cup \mathcal{U}\}$. Suppose $s \xrightarrow{\mathcal{P}} t$ or $s \xrightarrow{\mathcal{U}} t$ respectively, and thus $t = C[s_1^\#, \dots, t_i, \dots, s_n^\#]$ for some term t_i . There exists a context C' (over sequences) such that $\mathbf{N}_\pi^s(s) = C'[\mathbf{N}_\pi^s(s_i^\#)]$ and $\mathbf{N}_\pi^s(t) = C'[\mathbf{N}_\pi^s(t_i^\#)]$. First assume $s_i^\# \xrightarrow{\mathcal{P}} t_i$, and thus $\mathbf{N}_\pi^s(s_i^\#) = [\mathbf{N}_\pi(l^\#\sigma)]$ and $\mathbf{N}_\pi^s(t_i) = [[\mathbf{N}_\pi(r_1^\#\sigma)], \dots, [\mathbf{N}_\pi(r_m^\#\sigma)]]$ for $l \rightarrow c(r_1^\#, \dots, r_m^\#) \in \mathcal{P}$. To verify $\mathbf{N}_\pi^s(s) \blacktriangleright_k \mathbf{N}_\pi^s(t)$, by Definition 2.2(ii) and Definition 2.2(iv), it suffices to verify $\mathbf{N}_\pi(l^\#\sigma) \blacktriangleright_{k-1} \mathbf{N}_\pi(r_j^\#\sigma)$ for all $j \in \{1, \dots, m\}$. The latter is an easy consequence of Lemma 5.6, where we employ that (i) $l^\# >_{\text{pop}^*}^\pi r_j^\#$ follows from the assumption $\mathcal{P} \subseteq >_{\text{pop}^*}^\pi$, and (ii) $\|\pi(r)\| > \|\pi(r_j)\|$. Both properties are straight forward to verify since π is safe. For $s_i^\# \xrightarrow{\mathcal{U}} t$ we have $\mathbf{N}_\pi^s(s_i^\#) = [\mathbf{N}_\pi(s_i^\#)]$ and $\mathbf{N}_\pi^s(t_i^\#) = [\mathbf{N}_\pi(t_i^\#)]$ for $l \rightarrow r \in \mathcal{U}$. From Lemma 5.6 we obtain $\mathbf{N}_\pi(s_i^\#) \blacktriangleright_k \mathbf{N}_\pi(t_i^\#)$ which establishes the lemma. $\square$

The proof of Theorem 5.11 is now easily obtained by incorporating the above lemma into Theorem 5.9.

Theorem. Let $\mathcal{R}$ be a constructor TRS, and let $\mathcal{P}$ denote the set of weak innermost dependency pairs. Assume $\mathcal{P}$ is non-duplicating, and suppose $\mathcal{U}(\mathcal{P}) \subseteq >_{\mathcal{A}}$ for some SLI $\mathcal{A}$. Let π be a safe argument filtering. If $\mathcal{P} \subseteq >_{\text{pop}^*}^\pi$ and $\mathcal{U}(\mathcal{P}) \subseteq \gtrsim_{\text{pop}^*}^\pi$ then $\text{rc}_{\mathcal{R}}^i$ is polynomially bounded.

Proof. According to Proposition 3.4 we need to find a polynomial p such that

$$\text{dl}(t^\sharp, \xrightarrow{\text{WIDP}(\mathcal{R})/\mathcal{U}(\text{WIDP}(\mathcal{R}))}) \leq p(|t^\sharp|) .$$

We set $\mathcal{P} = \text{TPWIDP}(\mathcal{R})$ and likewise $\mathcal{U} = \mathcal{U}(\text{WIDP}(\mathcal{R}))$. Clearly, it suffices to show $\text{dl}(t^\sharp, \xrightarrow{\text{TPWIDP}(\mathcal{R})/\mathcal{U}(\text{WIDP}(\mathcal{R}))}) \leq p(|t^\sharp|)$ for that. Consider a sequence

$$t^\sharp = t_0 \xrightarrow{\mathcal{P}/\mathcal{U}} t_1 \xrightarrow{\mathcal{P}/\mathcal{U}} \dots \xrightarrow{\mathcal{P}/\mathcal{U}} t_\ell ,$$

and pick a relative step $t_i \xrightarrow{\mathcal{P}/\mathcal{U}} t_{i+1}$. Define $\mathcal{U}' = \mathcal{U} \cup \mathcal{V}(\mathcal{P} \cup \mathcal{U})$ and $\phi(t) = \phi_{\mathcal{P} \cup \mathcal{U}}(t)$. Clearly Lemma 5.7 can be extended to account for steps of $\mathcal{P}$ below the root, and thus $\phi(t_i) \xrightarrow{\mathcal{P}/\mathcal{U}'} \phi(t_{i+1})$ follows. Hence for some terms u and v , $\phi(t_i) \xrightarrow{\mathcal{U}'}^* u \xrightarrow{\mathcal{P}} v \xrightarrow{\mathcal{U}'}^* \phi(t_{i+1})$. As shown in Lemma A.10, all involved terms in the above sequence have the shape $C[s_1^\sharp, \dots, s_n^\sharp]$, $C \in \text{ComCtx}$. As $\text{WIDP}(\mathcal{R}) \subseteq >_{\text{pop}^*}^\pi$, and since π is safe, it is easy to infer that $\mathcal{P} \subseteq >_{\text{pop}^*}^\pi$ holds (we just set every compound symbol from $\mathcal{P}$ minimal in the precedence). And hence Lemma A.11 translates the above relative step to $\mathbf{N}_\pi^s(\phi(s)) \blacktriangleright_k^+ \mathbf{N}_\pi^s(\phi(t))$ for some uniform constant k . As a consequence, $\text{dl}(t, \xrightarrow{\text{WIDP}(\mathcal{R})/\mathcal{U}(\text{WIDP}(\mathcal{R}))}) \leq \mathbf{G}_k(\mathbf{N}_\pi^s(\phi(t)))$ for all terms t . Fix some reducible and basic term $t \in \mathcal{T}_{\mathbf{b}}$. Observe $\mathbf{N}_\pi^s(\phi(t^\sharp)) = [\mathbf{N}_\pi(t^\sharp)]$ and so from Lemma 5.1 we see that $\mathbf{G}_k(\mathbf{N}_\pi^s(\phi(t^\sharp)))$ is bounded polynomially in the size of t . The polynomial depends only on k . We conclude the theorem. $\square$

A.2. Proof of Theorem 5.14

We now proceed with the proof Theorem 5.14, which is essentially an extension to Theorem 5.11.

We first precisely state what it means that a TRS *computes* some function. For this, let $\ulcorner \cdot \urcorner : \Sigma^* \rightarrow \mathcal{T}(\mathcal{C})$ denote an *encoding function* that represents words over the alphabet Σ as ground values. We call an encoding $\ulcorner \cdot \urcorner$ *reasonable* if it is bijective and there exists a constant c such that $|u| \leq |\ulcorner u \urcorner| \leq c \cdot |u|$ for every $u \in \Sigma^*$. Let $\ulcorner \cdot \urcorner$ denote a reasonable encoding function, and let $\mathcal{R}$ be a completely defined, orthogonal and terminating TRS. We say that an n -ary function $f : (\Sigma^*)^n \rightarrow \Sigma^*$ is *computable* by $\mathcal{R}$ if there exists a defined function symbol $\mathbf{f}$ such that for all $w_1, \dots, w_n, v \in \Sigma^*$ $\mathbf{f}(\ulcorner w_1 \urcorner, \dots, \ulcorner w_n \urcorner) \rightarrow^! \ulcorner v \urcorner \iff f(w_1, \dots, w_n) = v$. On the other hand the TRS $\mathcal{R}$ *computes* f , if the function $f : (\Sigma^*)^n \rightarrow \Sigma^*$ is defined by the above equation.

Below we abbreviate $\mathbf{Q}_\pi$ as $\mathbf{Q}$ for predicative interpretation $\mathbf{Q} \in \{\mathbf{S}, \mathbf{N}, \mathbf{N}^s\}$ and the particular argument filtering π that induces the identity function on terms. Consider the following lemma.

Lemma A.12. *Let $\mathcal{R}$ be an S -sorted and completely defined constructor TRS such that the underlying signature is simple. If $\text{TPWIDP}(\mathcal{R}) \cup \mathcal{U}(\text{WIDP}(\mathcal{R})) \subseteq \succ_{\text{pop}^*}$ then there exists a polynomial p such that for all ground and well-typed basic terms $t \in \mathcal{T}_{\mathbf{b}}$, $t^\sharp \xrightarrow{\text{TPWIDP}(\mathcal{R}) \cup \mathcal{U}(\text{WIDP}(\mathcal{R}))}^* s$ implies $|s| \leq p(|t|)$.*

Proof. Let $\mathcal{S} = \text{TPWIDP}(\mathcal{R}) \cup \mathcal{U}(\text{WIDP}(\mathcal{R}))$. Suppose $t^\sharp \xrightarrow{\mathcal{S}}^* s$, or equivalently $t^\sharp \xrightarrow{\mathcal{S}}^* s$ since $\mathcal{R}$ is completely defined. By Lemma A.11 we derive $\mathbf{N}^s(t^\sharp) \blacktriangleright_k^* \mathbf{N}^s(s)$ for some uniform $k \in \mathbb{N}$. And thus $\mathbf{G}_k(\mathbf{N}^s(s)) \leq \mathbf{G}_k(\mathbf{N}^s(t^\sharp))$. As $\mathbf{G}_k(\mathbf{N}^s(t^\sharp)) = \mathbf{G}_k([\mathbf{N}(t^\sharp)])$ is bounded polynomially in the size of t according to Lemma 5.2, we see that there exists a polynomial p such that $\mathbf{G}_k(\mathbf{N}^s(s)) \leq \mathbf{G}_k(\mathbf{N}^s(t^\sharp)) \leq p(|t|)$. Since $\mathcal{R}$ is and $\mathcal{S}$ -sorted TRS over a simple signature, the same holds for $\mathcal{S}$ due to Lemma A.7. And thus since $t^\sharp$ is well-typed and $t^\sharp \xrightarrow{\mathcal{S}}^* s$ holds, also s is well-typed. Let q be the polynomial as given from Lemma A.4 with $|s| \leq q(\mathbf{G}_k(\mathbf{N}^s(s)))$. Summing up, we derive $|s| \leq q(\mathbf{G}_k(\mathbf{N}^s(s))) \leq q(p(|t|))$ as desired. $\square$

The above lemma has established that sizes of reducts with respect to the relation $\xrightarrow{\text{TPWIDP}(\mathcal{R}) \cup \mathcal{U}(\text{WIDP}(\mathcal{R}))}$ are bounded polynomially in the size of the start term, provided we can orient dependency pairs and usable rules. It remains to verify that this is indeed sufficient to appropriately estimate sizes of reducts with respect to $\xrightarrow{\mathcal{R}}$. The fact is established in the final Theorem.

Theorem. Let $\mathcal{R}$ be an orthogonal $\mathcal{S}$ -sorted and completely defined constructor TRS such that the underlying signature is simple. Let $\mathcal{P}$ denote the set of weak innermost dependency pairs. Assume $\mathcal{P}$ is non-duplicating, and suppose $\mathcal{U}(\mathcal{P}) \subseteq >_{\mathcal{A}}$ for some SLI $\mathcal{A}$. If $\mathcal{P} \subseteq >_{\text{pop}^*}^\pi$ and $\mathcal{U}(\mathcal{P}) \subseteq \gtrsim_{\text{pop}^*}^\pi$ then the functions computed by $\mathcal{R}$ are computable in polynomial-time.

Proof. We single out one of the defined symbols $\mathbf{f} \in \mathcal{D}$ and consider the corresponding function $f: (\Sigma^*)^n \rightarrow \Sigma^*$ computed by $\mathcal{R}$. Under the assumptions, $\mathcal{R}$ is terminating, but moreover $\text{rc}_{\mathcal{R}}^i$ is polynomially bounded according to Theorem 5.11. Additionally, from orthogonality (and hence confluence) of $\mathcal{R}$, normal forms are unique and so the function f is well-defined. Suppose $\mathbf{f}(\ulcorner w_1 \urcorner, \dots, \ulcorner w_n \urcorner) \xrightarrow{\mathcal{R}}^! \ulcorner v \urcorner$ for words $w_1, \dots, w_n, v$. In particular, from confluence we see that

$$\mathbf{f}(\ulcorner w_1 \urcorner, \dots, \ulcorner w_n \urcorner) \xrightarrow{\mathcal{R}} t_1 \xrightarrow{\mathcal{R}} \dots \xrightarrow{\mathcal{R}} t_\ell = \ulcorner v \urcorner.$$

It is folklore that there exists a polytime algorithm performing one rewrite step. Hence to conclude the existence of a polytime algorithm for f it suffices to bound the size of terms t_i for $1 \leq i \leq \ell$ polynomially in $\sum_i |w_i|$. And as we suppose that the encoding $\ulcorner \cdot \urcorner$ is reasonable, it thus suffice to bound the sizes of t_i for $i \in \{1, \dots, \ell\}$ polynomially in the size of $t_0 = \mathbf{f}(\ulcorner w_1 \urcorner, \dots, \ulcorner w_n \urcorner)$.

Consider a term t_i . Without loss of generality, we can assume t_i is ground. According to Lemma A.10 there exists contexts $C'_i \in \mathbf{ComCtx}$, $C_i \in \mathbf{reprs}(C'_i)$ and terms $u_1, \dots, u_n$ such that $t_i = C_i[u_1, \dots, u_n]$ and moreover, $t_0^\sharp \xrightarrow{\mathcal{P} \cup \mathcal{U}}^* C'_i[u_1^\sharp, \dots, u_n^\sharp]$ for all $i \in \{1, \dots, \ell\}$. From the assumption $\text{WIDP}(\mathcal{R}) \subseteq >_{\text{pop}^*}$ we see $\text{TPWIDP}(\mathcal{R}) \subseteq >_{\text{pop}^*}$. Thus by Lemma A.12 there exists a polynomial p such that $|C'_i[u_1^\sharp, \dots, u_n^\sharp]| \leq p(|t_0|)$. And so, clearly $\sum_{j=0}^n |u_j| \leq p(|t_0|)$. It remains to bound the sizes of contexts C_i polynomially in $|t_0|$.

Recall Definition A.8, and recall that $C_i \in \mathbf{reprs}(C'_i)$. Thus C_i is a context build from constructors and variables, where the latter are replaced by normal forms of $\mathcal{R}$.

Since $\mathcal{R}$ is completely defined, $\mathbf{NF}(\mathcal{R})$ coincides with values. We conclude that $C_i \in \mathcal{T}(\mathcal{C} \cup \{\Box_s \mid s \in S\})$. Here $\Box_s$ denotes the hole of sort s . Moreover since $\mathcal{R}$ is S -sorted, and $t_0 \xrightarrow{\mathcal{R}}^* C_i[u_1, \dots, u_n]$, we see that C_i is well-typed. We define $\Delta_{\mathcal{R}} = \max\{\mathbf{dp}(r) \mid l \rightarrow r \in \mathcal{R}\}$. By a straight forward induction it follows that $\mathbf{dp}(t_i) \leq \mathbf{dp}(t_0) + \Delta_{\mathcal{R}} \cdot i \leq |t_0| + \Delta_{\mathcal{R}} \cdot \mathbf{dl}(t_0, \xrightarrow{\mathcal{R}})$. As a consequence, $\mathbf{dp}(C_i) \leq |t_0| + \Delta_{\mathcal{R}} \cdot \mathbf{dl}(t_0, \xrightarrow{\mathcal{R}})$, and thus by Proposition A.3 there exists constants $c, d \in \mathbb{N}$ such that $|C_i| \leq c \cdot \mathbf{dp}(C_i)^d \leq c \cdot (|t_0| + \Delta_{\mathcal{R}} \cdot \mathbf{dl}(t_0, \xrightarrow{\mathcal{R}}))^d$. As we have that $\mathbf{dl}(t_0, \xrightarrow{\mathcal{R}})$ is polynomially bounded in the size of t_0 , it follows that $|C_i| \leq q(|t_0|)$ for some polynomial q .

Summing up, we conclude that for all $i \in \{1, \dots, \ell\}$, $|t_i| \leq p(|t_0|) + q(|t_0|)$ for the polynomials p and q from above. This concludes the theorem. $\square$

References

- [1] Toshiyasu Arai and Georg Moser. Proofs of termination of rewrite systems for polytime functions. In *Proc. 25th FSTTCS*, volume 3821 of *LNCS*, pages 529–540, 2005.
- [2] Thomas Arts and Jürgen Giesl. Termination of term rewriting using dependency pairs. *TCS*, 236(1–2):133–178, 2000.
- [3] Martin Avanzini. Automation of polynomial path orders. Master’s thesis, University of Innsbruck, Faculty for Computer Science., 2009. Available at <http://cl-informatik.uibk.ac.at/~zini/MT.pdf>.
- [4] Martin Avanzini and Georg Moser. Complexity analysis by rewriting. In *Proc. 9th FLOPS*, volume 4989 of *LNCS*, pages 130–146, 2008.
- [5] Martin Avanzini, Georg Moser, and Andreas Schnabl. Automated implicit computational complexity analysis (system description). In *Proc. 4th IJCAR*, volume 5195 of *LNCS*, pages 132–138, 2008.
- [6] Franz Baader and Tobias Nipkow. *Term Rewriting and All That*. Cambridge University Press, 1998.
- [7] Stephen Bellantoni and Stephen Cook. A new recursion-theoretic characterization of the polytime functions. *CC*, 2(2):97–110, 1992.
- [8] G. Bonfante, J.-Y. Marion, and J.-Y. Moyen. Quasi-interpretations: A way to control resources. *TCS*, 2009. To appear.
- [9] Niklas Eén and Niklas Sörensson. An extensible SAT-solver. In *Proc. 6th SAT*, volume 2919 of *LNCS*, pages 502–518, 2003.
- [10] Jörg Endrullis, Johannes Waldmann, and Hans Zantema. Matrix interpretations for proving termination of term rewriting. *JAR*, 40(2–3):195–220, 2008.

- [11] Carsten Fuhs, Jürgen Giesl, Aart Middeldorp, Peter Schneider-Kamp, René Thiemann, and Harald Zankl. SAT solving for termination analysis with polynomial interpretations. In *Proc. 10th SAT*, volume 4501 of *LNCS*, pages 340–354, 2007.
- [12] Alfons Geser. *Relative Termination*. PhD thesis, University of Passau, Faculty for Mathematics and Computer Science, 1990.
- [13] Jürgen Giesl, René Thiemann, and Peter Schneider-Kamp. Proving and disproving termination of higher-order functions. In *Proc. 5th FroCoS*, volume 4501 of *LNCS*, pages 340–354, 2005.
- [14] Nao Hirokawa and Georg Moser. Complexity, graphs, and the dependency pair method. In *Proc. 15th LPAR*, volume 5330 of *LNCS*, pages 667–681, 2008.
- [15] Nao Hirokawa and Georg Moser. Automated complexity analysis based on the dependency pair method. In *Proc. 4th IJCAR*, volume 5195 of *LNCS*, pages 364–380, 2008.
- [16] D. Hofbauer and C. Lautemann. Termination proofs and the length of derivations. In *Proc. 3rd RTA*, volume 355 of *LNCS*, pages 167–177, 1989.
- [17] J.-Y. Marion and R. Péchoux. Characterizations of polynomial complexity classes with a better intensionality. In *Proc. 10th PPDP*, pages 79–88. ACM, 2008.
- [18] Jean-Yves Marion. Analysing the implicit complexity of programs. *IC*, 183:2–18, 2003.
- [19] Harold Simmons. The realm of primitive recursion. *ARCH*, 27:177–188, 1988.
- [20] Terese. *Term Rewriting Systems*, volume 55 of *Cambridge Tracts in Theoretical Computer Science*. Cambridge University Press, 2003.
- [21] René Thiemann. *The DP Framework for Proving Termination of Term Rewriting*. PhD thesis, University of Aachen, Department of Computer Science, 2007.